



Herbicide

ACCEPTED
FOR REGISTRATION

August 10, 2023

DOC ID# 588278

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration

Classified for
“RESTRICTED USE”
in New York State
under 6NYCRR Part 326



METSULFURON-METHYL GROUP 2 HERBICIDE

Dry Flowable	
Active Ingredient	By Weight
Metsulfuron methyl	
Methyl 2-[[[4-methoxy-6-methyl-1,3,5-triazin-2-yl]amino]-carbonyl]amino)sulfonyl]benzoate.....	60%
Other Ingredients	40%
TOTAL	100%
EPA Reg. No. 101563-167 EPA Est. No.	

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)
See Panel for First Aid Instructions and Booklet for Complete Precautionary Statements and Directions for Use.

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.
IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

85798669
85796941D 221216AV1
Nonrefillable Container
Net Weight:
1 Pound

PRODUCED FOR
Environmental Science U.S., LLC
5000 CentreGreen Way, Suite 400
Cary, NC 27513

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! Causes eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep dry and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

This herbicide is injurious to plants at extremely low concentrations. Nontarget plants may be adversely effected from drift and run-off.

GROUNDWATER ADVISORY

Metsulfuron-methyl is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

SURFACE WATER ADVISORY

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for weeks after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of metsulfuron-methyl from runoff water and sediment. Runoff of this product will be greatly reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

NON-TARGET ORGANISM ADVISORY

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by minimizing spray drift. For further guidance and instructions on how to minimize spray drift, refer to the Spray Drift Management section of this label.

WINDBLOWN SOIL PARTICLES

ESCORT® XP HERBICIDE has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying ESCORT® XP HERBICIDE if prevailing local conditions may be expected to result in off-site movement.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ESCORT® XP HERBICIDE must be used only in accordance with instructions on this label or in separately published ENVIRONMENTAL SCIENCE U.S., LLC instructions. ENVIRONMENTAL SCIENCE U.S., LLC will not be responsible for losses or damages resulting from the use of this product in any manner not specified on this label. User assumes all risks associated with such non-specified use.

Do not apply more than 4 ounces of ESCORT® XP HERBICIDE (0.15 pounds of the active ingredient metsulfuron-methyl) per acre per year.

Do not use on food or feed crops except as specified by this label or supplemental labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency in your State responsible for pesticide regulation.

PRODUCT INFORMATION

ESCORT® XP HERBICIDE is a dispersible granule that is mixed in water and applied as a spray by ground or aerial application.

ESCORT® XP HERBICIDE is registered for the control of annual and perennial weeds and unwanted woody plants on private, public and military lands, on rights-of-way, industrial sites, non-crop areas, ditchbanks of dry drainage ditches, certain types of unimproved turf grass, and conifer and hardwood plantations, including grazed areas on these sites. Do not use on irrigation ditches.

ESCORT® XP HERBICIDE controls weeds and woody plants primarily by postemergence activity. Although ESCORT® XP HERBICIDE has preemergence activity, best results are generally obtained when ESCORT® XP HERBICIDE is applied to foliage after emergence or dormancy break. Generally, for the control of annual weeds, ESCORT® XP HERBICIDE provides the best results when applied to young, actively growing weeds. For the control of perennial weeds, applications made at the bud/bloom stage or while the target weeds are in the fall rosette stage may provide the best results. The use rate depends upon the weed species and size at the time of application.

The degree and duration of control may depend on the following:

- weed spectrum and infestation intensity
- weed size at application
- environmental conditions at and following treatment
- soil pH, soil moisture, and soil organic matter

ESCORT® XP HERBICIDE may be applied on conifer and hardwood plantations, and non-crop sites that contain areas of temporary surface water caused by the collection of water between planting beds, in equipment ruts, or in other depressions created by management activities. It is permissible to treat intermittently flooded low lying sites, seasonally dry flood plains and transitional areas between upland and lowland sites when no water is present. It is also permissible to treat marshes, swamps and bogs after water has receded as well as seasonally dry flood deltas. DO NOT make applications to natural or man-made bodies of water such as lakes, reservoirs, ponds, streams, and canals.

BIOLOGICAL ACTIVITY

ESCORT® XP HERBICIDE is absorbed primarily through the foliage of plants, and by the roots to a lesser degree. Plant cell division is generally inhibited in sensitive plants within a few hours following uptake. Two to 4 weeks after application, leaf growth slows followed by discoloration and tissue death. The final effects on annual weeds are evident about 4 to 6 weeks after application. The ultimate affect on perennial weeds and woody plants occurs in the growing season following application.

Warm, moist conditions following treatment promote the activity of ESCORT® XP HERBICIDE, while cold, dry conditions may reduce or delay activity. Weeds and brush hardened off by cold weather or drought stress may not be controlled. Weed and brush control may be reduced if rainfall occurs soon after application.

ADJUVANTS

The use of a surfactant is recommended to enhance the control of susceptible plants, except where noted. Apply at a minimum rate (concentration) of 1/4% volume/volume (1 quart per 100 gallons of spray solution), or at the manufacturer's recommended rate. Use only EPA approved surfactants containing at least 80% active ingredient. Certain types of surfactants, such as those incorporating acetic acid (i.e. LI-700), may not be compatible with ESCORT® XP HERBICIDE and may result in decreased performance. Certain surfactants may not be suitable for use on desirable plants, such as turf and conifers, listed on this label. Consult the surfactant manufacturer's label for appropriate uses.

INVASIVE SPECIES MANAGEMENT

This product may be considered for use on public, private, and tribal lands to treat certain weed species infestations that have been determined to be invasive, consistent with the Federal Interagency Committee for the Management of Noxious and Exotic Weeds (FICMNEW) National Early Detection and Rapid Response (EDRR) System for invasive plants.

Effective EDRR systems address invasions by eradicating the invader where possible, and controlling them when the invasive species is too established to be feasibly eradicated. Once an EDRR assessment has been completed and action is recommended, a Rapid Response needs to be taken to quickly contain, deny reproduction, and if possible eliminate the invader. Consult your appropriate state extension service, forest service, or regional multidisciplinary invasive species management coordination team to determine the appropriate Rapid Response.

WEED RESISTANCE MANAGEMENT

ESCORT® XP HERBICIDE contains the active ingredient metsulfuron-methyl which is a Group 2 Herbicide based on the mode of action classification system of the Weed Science Society of America. When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected.

Follow the best management practices listed below to delay the development of herbicide resistant weeds.

- Fields should be scouted prior to application to identify the weed species present and their growth stage to determine if the intended application will be effective. Fields should be scouted after application to verify that the treatment was effective.
- Identify weeds present in the field through scouting and field history and understand their biology. The weed-control program should consider all of the weeds present.
- Suspected herbicide-resistant weeds may be identified by these indicators:
 - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - A spreading patch of non-controlled plants of a particular weed species; and
 - Surviving plants mixed with controlled individuals of the same species.
- Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of actions for each target weed.
- Report any incidence of non-performance of this product against a particular weed species to your Environmental Science U.S., LLC distributor, Environmental Science U.S., LLC representative or call 1-800-331-2867.
- If resistance is suspected, treat weed escapes with an herbicide having a different mechanism of action and/or use non-chemical means to remove escapes, as practical, with the goal of preventing further seed production.
- Use a diversified approach toward weed management. Whenever possible incorporate multiple weed-control practices such as mechanical cultivation, biological management practices, and crop rotation.
- To the extent possible, do not allow weed escapes to produce seeds, roots, or tubers.
- Difficult to control weeds may require sequential applications of herbicides with differing mechanisms of action.

- Apply this herbicide at the correct timing and rate needed to control the most difficult weeds in the field.
- Use a broad spectrum soil-applied herbicide with a mechanism of action that differs from this product as a foundation in a weed-control program.
- Do not use more than two applications of this or any other herbicide with the same mechanism of action within a single growing season unless mixed with an herbicide with another mechanism of action with an overlapping spectrum for the difficult-to-control weeds.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants, or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

PREPARING FOR USE - Site Specific Considerations

Understanding the risks associated with the application of ESCORT® XP HERBICIDE is essential to aid in preventing off-site injury to desirable vegetation and agricultural crops. The risk of off-site movement, both during and after application, may be affected by a number of site specific factors such as the nature, texture and stability of the soil, the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, drainage patterns, and other local physical and environmental conditions. A careful evaluation of the potential for off-site movement from the intended application site, including movement of treated soil by wind or water erosion, must be made prior to using ESCORT® XP HERBICIDE. This evaluation is particularly critical where desirable vegetation or crops are grown on neighboring land for which the use of ESCORT® XP HERBICIDE is not labeled. If prevailing local conditions may be expected to result in off-site movement and cause damage to neighboring desirable vegetation or agricultural crops, do not apply ESCORT® XP HERBICIDE.

Before applying ESCORT® XP HERBICIDE the user must read and understand all label directions, precautions and restrictions completely, including these requirements for a site specific evaluation. If you do not understand any of the instructions or precautions on the label, or are unable to make a site specific evaluation yourself, consult your local agricultural dealer, cooperative extension service, land managers, professional consultants, or other qualified authorities familiar with the area to be treated. If you still have questions regarding the need for site specific considerations, please call 1-800-331-2867.

TANK MIXES

ESCORT® XP HERBICIDE may be tank mixed with other herbicides registered for the use sites described in this label. Use only those tank mix partners which are labeled for the appropriate use site. When tank mixing, use the most restrictive label limitations for each of the products being used in the tank mix.

AGRICULTURAL USES

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Shoes plus socks

CONIFER PLANTATIONS

Application Information

ESCORT® XP HERBICIDE is registered for the control of many species of weeds and deciduous trees on sites where conifers are growing or are to be planted. Apply by ground equipment or by air (helicopter only). Refer to the "Weeds Controlled" and "Brush Species Controlled" for a listing of susceptible species.

Application Timing

Apply ESCORT® XP HERBICIDE after weeds have emerged or after undesirable hardwoods have broken winter dormancy and have reached the point of full leaf expansion.

Conifer Site Preparation

--Application Before Transplanting

After consulting the "Weeds Controlled" and "Brush Species Controlled" tables, apply the rates of ESCORT® XP HERBICIDE specified for the most difficult to control species on the site.

Southeast—Apply up to 4 ounces per acre for loblolly and slash pines. Transplant the following planting season.

Northeast and Lake States—Apply up to 2 ounces per acre for red pine. Transplant the following planting season. Apply up to 2 ounces per acre for black, white and Norway spruce. Transplant the following spring.

West—Apply up to 2 ounces per acre prior to planting Douglas Fir, Sitka Spruce, Western Red Cedar, Western Hemlock, Ponderosa Pine, and Grand Fir in the Coast Rangeland and western slope of the Cascades in Oregon and Washington. These conifer species listed can be planted anytime after application. Other conifer species can be planted providing the user has prior experience indicating acceptable tolerance to ESCORT® XP HERBICIDE soil residues. Without prior experience, it is recommended that other species be planted on a small scale to determine selectivity before large-scale plantings are made as unacceptable injury may occur. ENVIRONMENTAL SCIENCE U.S., LLC will not assume responsibility for injury to any conifer species not listed on this label.

Tank Mix Combinations—

For broader spectrum control, the following products may be used in combination with ESCORT® XP HERBICIDE.

Glyphosate (4 pound active per gallon)

Tank mix 1 to 2 ounces of ESCORT® XP HERBICIDE with 2 to 10 quarts of glyphosate per acre. Refer to the product container for a list of species controlled.

Imazapyr (4 pound active per gallon)

Tank mix 1 to 2 ounces of ESCORT® XP HERBICIDE with 10 to 24 fluid ounces of imazapyr per acre. Loblolly and slash pines may be transplanted the planting season following application. This combination controls ash, black gum, cherry, hawthorn, honeysuckle, hophornbeam, persimmon, oaks (red, white and water), sassafras, sweetgum, Vaccinium species, and suppresses blackberry, dogwood, elms, myrtle dahoon, hickories, and red maple.

Glyphosate (4 pound active per gallon) + Imazapyr (4 pound active per gallon)

Tank mix 1/2 to 1 ounce of ESCORT® XP HERBICIDE with 16 to 64 fluid ounces of glyphosate and 10 to 12 fluid ounces of imazapyr per acre. Slash and loblolly pines may be transplanted the planting season following application. This combination controls cherry, dogwood, elms, oaks (red and water), persimmon, sassafras, sweetgum and suppresses hickory.

VELPAR® L VU HERBICIDE or VELPAR® DF VU HERBICIDE

Tank mix 1 to 2 ounces of ESCORT® XP HERBICIDE per acre with VELPAR® L VU HERBICIDE or VELPAR® DF VU HERBICIDE at the rates specified on the container for various soil textures. Loblolly and slash pines may be transplanted the planting season following application. Refer to the product container for a list of species controlled.

OUST® EXTRA HERBICIDE

Tank mix 1/2 to 1 1/2 ounces of ESCORT® XP HERBICIDE with 2 to 3 ounces of OUST® EXTRA HERBICIDE per acre for herbaceous weed control. Refer to the product container and the "Weeds Controlled" section of this label for a listing of the weeds controlled. Loblolly and slash pines may be transplanted the planting season following application. Tank mix 2 ounces of ESCORT® XP HERBICIDE with 3 ounces of OUST® EXTRA HERBICIDE per acre for herbaceous weed control and early spring suppression of bull thistle and Canada thistle in the Coast Rangeland and western slope of the Cascade Mountains. Douglas fir may be transplanted at least 90 days following application.

Release--Hardwood Control and Suppression

ESCORT® XP HERBICIDE may be used for application over the top of established slash and loblolly pine to control the species listed in "Weeds Controlled" and "Brush Species Controlled" section of this label. Apply 1 to 4 ounces per acre to control the species indicated, including kudu.

Tank Mix Combinations—

For broader spectrum control the following products may be used in combination with ESCORT® XP HERBICIDE.

Imazapyr (4 pound active per gallon)

Tank mix 1 to 2 ounces of ESCORT® XP HERBICIDE with 8 to 16 fluid ounces of imazapyr per acre for application to loblolly pine. Refer to the imazapyr label regarding the use of surfactants and the appropriate application timing with respect to the age and development stage of the pines. This combination controls ash, black gum, cherry, hawthorn, honeysuckle, hophornbeam, oaks (red, white and water), sassafras, sweetgum, Vaccinium species, and suppresses blackberry, dogwood, elms, myrtle dahoon, hickories, persimmon, and red maple.

VELPAR® L VU HERBICIDE or VELPAR® DF VU HERBICIDE

Tank mix 1 to 2 ounces of ESCORT® XP HERBICIDE with VELPAR® L VU HERBICIDE or VELPAR® DF VU HERBICIDE at the rates specified on the container for various soil textures. This combination may be applied to loblolly and slash pines.

Release--Herbaceous Weed Control

ESCORT® XP HERBICIDE may be applied to transplanted loblolly and slash pine for the control of herbaceous competition. Consult the "Weeds Controlled" for a listing of the susceptible species and application rates. Best results are obtained when ESCORT® XP HERBICIDE is applied just before weed emergence until shortly after weed emergence.

Tank Mix Combinations—

For broader spectrum control the following products may be used in combination with ESCORT® XP HERBICIDE.

Imazapyr (4 pound active per gallon)

Tank mix 1/2 to 1 ounce of ESCORT® XP HERBICIDE with 4 fluid ounces of imazapyr per acre. The tank mix may be used on loblolly pine.

VELPAR® L VU HERBICIDE or VELPAR® DF VU HERBICIDE

Tank mix 1/2 to 1 ounce of ESCORT® XP HERBICIDE with VELPAR® L VU HERBICIDE or VELPAR® DF VU HERBICIDE at the rates specified on the container for various soil textures. This combination may be applied to loblolly and slash pines.

Release - Directed Spray in Conifers

Western US

To release conifers from competing brush species, such as, blackberry, salmonberry, snowberry, thimbleberry and wild roses, mix 2 to 4 ounces of ESCORT® XP HERBICIDE per 100 gallons of spray solution. Direct spray onto the foliage of competing brush species using a knapsack or backpack sprayer. For best results, apply any time after the brush species have reached full leaf stage but before autumn coloration. For best results at application, the majority of the brush must be less than six feet in height to help ensure adequate spray coverage. Thorough coverage of the target foliage is necessary to optimize results. Care must be taken to direct the ESCORT® XP HERBICIDE spray solution away from the conifer foliage.

NOTE:

ESCORT® XP HERBICIDE may cause temporary yellowing and or growth suppression when the spray solution contacts conifer foliage. The use of a surfactant with ESCORT® XP HERBICIDE may improve brush control results. When using a surfactant with ESCORT® XP HERBICIDE, extra precaution must be taken to avoid contact with conifer foliage. Excessive drift onto conifers may result in severe injury.

IMPORTANT PRECAUTIONS—CONIFER PLANTATIONS ONLY

- Applications of ESCORT® XP HERBICIDE made to conifers that are suffering from loss of vigor caused by insects, diseases, drought, winter damage, animal damage, excessive soil moisture, planting shock, or other stresses may injure or kill the trees.

- Applications of ESCORT® XP HERBICIDE made for herbaceous release must only be made after adequate rainfall has closed the planting slit and settled the soil around the roots following transplanting.
- Do not apply ESCORT® XP HERBICIDE to conifers grown as ornamentals.
- ESCORT® XP HERBICIDE applications may result in damage and mortality to other species of conifers when they are present on sites with those listed in the preceding specifications for conifer plantations.

HARDWOOD PLANTATIONS

Application Information

ESCORT® XP HERBICIDE may be used at rates of up to 2 ounces per acre for the control of many weed species on sites where yellow poplar is growing or is to be planted, and on sites where red alder is to be planted. Apply by ground equipment or by air (helicopter only). Refer to the "Weeds Controlled" sections of this label for a listing of susceptible species.

Application Timing

ESCORT® XP HERBICIDE may be applied as a site preparation treatment prior to planting red alder or yellow poplar. As a prior to planting site preparation treatment for red alder, ESCORT® XP HERBICIDE may be tank mixed with other herbicides labeled for this use.

ESCORT® XP HERBICIDE may also be applied over-the-top of planted yellow poplar seedlings after the soil has settled around the root system, but before the seedlings have broken dormancy (prior to bud-break).

Release--Herbaceous Weed Control

ESCORT® XP HERBICIDE may be applied to yellow poplar for the control of herbaceous competition. Consult the "Weeds Controlled" for a listing of the susceptible species and specified application rates. Best results are obtained when ESCORT® XP HERBICIDE is applied just before weed emergence until shortly after weed emergence.

Tank Mix Combinations—

Tank mix 1/2 ounce of ESCORT® XP HERBICIDE with 4 to 6 pints of VELPAR® L VU HERBICIDE as directed on the package label for "RELEASE--HERBACEOUS WEED CONTROL" in pine plantations in the eastern U.S. Follow the VELPAR® L VU HERBICIDE label directions regarding altering the application rate by soil texture.

IMPORTANT PRECAUTIONS—HARDWOOD PLANTATIONS ONLY

- Application of VELPAR® L VU HERBICIDE and ESCORT® XP HERBICIDE made to yellow poplar that are suffering from loss of vigor caused by insects, disease, drought, winter damage, animal damage, excessive soil moisture, planting shock, or other stresses may injure or kill the seedlings.
- Applications of ESCORT® XP HERBICIDE made for release must only be made after adequate rainfall has closed the planting slit and settled the soil around the roots following transplanting.
- The use of surfactant is not recommended for applications made over the tops of trees.
- Careful consideration must be given by an experienced and knowledgeable forester to match the requirements of yellow poplar and/or red alder to the conditions of the site. Treatment of yellow poplar and/or red alder planted on a site inadequate to meet its requirements may injure or kill the seedlings.

PASTURE, RANGELAND, AND CONSERVATION RESERVE PROGRAM (CRP)

ESCORT® XP HERBICIDE is registered for the control of broadleaf weeds, brush and several woody vine species in the establishment, maintenance, and restoration of pasture, rangeland, and Conservation Reserve Program (CRP).

ESCORT® XP HERBICIDE may be tank mixed with other pesticides labeled for use in pasture, rangeland, and CRP. Read and follow the labels on all products used in the tank mix. Observe the most restrictive precautions on each of the product's labels. Application of ESCORT® XP HERBICIDE to pasture, rangeland, and CRP may be made by ground or air. Use a sufficient volume of water to ensure thorough coverage of the targeted weeds with the equipment being used. In Idaho, Oregon and Washington use a minimum application volume of 3 gallons of spray solution per acre.

APPLICATION INFORMATION FOR GRASS ESTABLISHMENT IN PASTURE, RANGELAND, AND CONSERVATION RESERVE PROGRAM (CRP)

ESCORT® XP HERBICIDE is registered for the control or suppression of broadleaf weeds to aid in the establishment of the following perennial native or improved grasses planted in pasture, rangeland, and acres enrolled in the Conservation Reserve Program (CRP):

Blue Gramma	Lovegrasses-	Wheatgrasses-	Wildrye grass-
Bluestems-	Atherstone	bluebunch	Russian
big	sand	crested	
little	weeping	intermediate	
plains	Wilman	pubescent	
sand	Orchardgrass	Siberian	
WW Spar	Sideoats gramma	slender	
Buffalograss	Switchgrass-	steambank	
Green spratletop	Blackwell	tall	
Kleingrass		thickspike	
		western	

Maximize potential for grass establishment by consulting with the Natural Resource and Conservation Service of other government agencies or local experts concerning planting techniques and other cultural practices.

Performance from ESCORT® XP HERBICIDE may not always be satisfactory due to the inability of newly planted grass stands to sufficiently compete with weeds and the severity of weed pressure in new grass stands.

An additional herbicide application or mowing may be needed.

Use Rates and Application Timing for Grass Establishment in Pasture, Rangeland, and CRP

Preplant (prior to planting) or Preemergence (after planting but before grass emergence)

Do not use more than 1/10 ounce/acre of ESCORT® XP HERBICIDE for grass establishment in pasture, rangeland, and CRP.

Apply ESCORT® XP HERBICIDE at 1/10 ounce/acre on all labeled grasses except orchardgrass and Russian wildrye grass. Do not apply ESCORT® XP HERBICIDE preplant or preemergence to orchardgrass and Russian wildrye grass as severe crop injury may result.

Early postemergence to new plantings

Apply ESCORT® XP HERBICIDE at 1/10 ounce/acre, plus a non-ionic surfactant at the rate of 2 to 4 pints/100 gallons of spray solution on all labeled grasses anytime after grass emergence.

Do not use a spray adjuvant other than non-ionic surfactant. Because grass species differ in time of emergence, apply only after the majority of grasses are in the 3 to 4 leaf stage.

Postemergence to stands with 1 – 5 leaf grasses planted the previous season.

Apply ESCORT® XP HERBICIDE at 1/10 ounce/acre plus a non-ionic surfactant at the rate of 2 to 4 pints/100 gallons of spray solution on all labeled grasses when the majority of the grasses have one or more leaves.

Do not use a spray adjuvant other than non-ionic surfactant.

APPLICATION INFORMATION FOR ESTABLISHED GRASSES IN PASTURE, RANGELAND, AND CONSERVATION RESERVE PROGRAM (CRP)

Use Rates for Established Grasses in Pasture, Rangeland, and CRP

Apply up to 1 2/3 ounces ESCORT® XP HERBICIDE per acre as a broadcast application to established grasses in pasture, rangeland, and CRP. For spot applications, use 1 ounce per 100 gallons of water. Do not apply more than 1 2/3 ounces of ESCORT® XP HERBICIDE per acre per year in pasture, rangeland, and CRP.

Refer to the Weeds Controlled section of the section 3 label for a listing of the weeds controlled by ESCORT® XP HERBICIDE and the appropriate use rate to obtain control.

Application Timing – Established Grasses in Pasture, Rangeland, and CRP

ESCORT® XP HERBICIDE may be applied to established native grasses such as bluestems and grama, and on other established grasses such as bermudagrass, bluegrass, orchardgrass, bromegrass, fescue and timothy that were planted the previous growing season (or earlier) and are fully tillered, unless otherwise directed on this label. Specific application timing information on several of these grass species follows:

Grass	Minimum time from Grass establishment ESCORT® XP HERBICIDE application
Bermudagrass	2 months
Bluegrass, bromegrass, Orchardgrass	6 months
Timothy	12 months
Fescue	24 months

Rotation Intervals in Pasture, Rangeland, and CRP for Overseeding and Renovation

Location	Crop or Grass Species	Maximum ESCORT® XP HERBICIDE Rate on Pasture, Rangeland, and CRP (oz per A)	Minimum Rotation Interval (months)
AL, AR, FL, GA, KY, LA, MS, NC, OK, SC, TN, TX, VA, WV	Alfalfa, red clover, white clover, sweet clover, bermudagrass, bluegrass, ryegrass, tall fescue	1/10 to 3/10	4
	Wheat (except durum)	1/10 to 3/10	1
	Durum, barley, oat	1/10 to 3/10	10
ALL STATES NOT INCLUDED ABOVE	Red clover, white clover, and sweet clover	1/10 to 2/10	12
	Bermudagrass, bluegrass, ryegrass	1/10 to 2/10	6
	Tall Fescue	1/10 to 2/10	18
	Wheat (except durum)	1/10 to 2/10	1
	Durum, barley, oat	1/10 to 2/10	10

(continued)

Rotation Intervals in Pasture, Rangeland, and CRP for Overseeding and Renovation (continued)

Location	Crop or Grass Species	Maximum ESCORT® XP HERBICIDE Rate on Pasture, Rangeland, and CRP (oz per A)	Minimum Rotation Interval (months)
ALL AREAS WITH SOIL PH OF 7.5 OR LESS	Russian wildrye	1/10 to 1/2	1
	Green needlegrass, switchgrass, sheep fescue	1/10 to 1	1
	Meadow brome, smooth brome, alta fescue, red fescue, meadow foxtail, orchardgrass, Russian wildrye, timothy	1/10 to 1	2
ALL AREAS WITH SOIL PH OF 7.9 OR LESS	Alkali sacaton, mountain brome, blue grama, thickspike wheatgrass	1/10 to 1	1
	Sideoats grama, switchgrass	1/10 to 1/2	2
	Western wheatgrass	1/10 to 1	2
	Sideoats grama, switchgrass, big bluestem	1/10 to 1	3

Fescue Precautions:

Note that ESCORT® XP HERBICIDE may temporarily stunt tall fescue, cause it to turn yellow, or cause seedhead suppression. To minimize these symptoms, take the following precautions:

- Do not use more than 4/10 ounce/acre of ESCORT® XP HERBICIDE
- Tank mix ESCORT® XP HERBICIDE with 2,4-D
- Use the lowest specified rate for target weeds

- Use a non-ionic surfactant at 1/2 to 1 pint per 100 gallons of spray solution
- Make application later in the spring after the new growth is 5 to 6 inches tall, or in the fall
- Do not use surfactant when liquid nitrogen is used as a carrier
- Do not use a spray adjuvant other than non-ionic surfactant

The first cutting yields may be reduced due to seedhead suppression resulting from treatment with ESCORT® XP HERBICIDE.

Timothy Precautions:

Timothy should be at least 6 inches tall at application and be actively growing. Applications of ESCORT® XP HERBICIDE to timothy under any other conditions may cause crop yellowing and/or stunting. To minimize these symptoms, take the following precautions:

- Do not use more than 4/10 ounce/acre ESCORT® XP HERBICIDE
- Tank mix ESCORT® XP HERBICIDE with 2, 4-D
- Use the lowest specified rate for target weeds
- Use a non-ionic surfactant at 1/2 pint per 100 gallons of spray solution (1/16%)
- Make applications in the late summer or fall
- Do not use surfactant when liquid nitrogen is used as a carrier
- Do not use spray adjuvant other than non-ionic surfactant

Application of ESCORT® XP HERBICIDE to Pensacola bahiagrass, ryegrass (Italian or perennial) and Garrison's creeping foxtail may cause severe injury to and/or loss of forage.

Other Pasture and Rangeland Grasses

Varieties and species of forage grasses differ in their tolerance to herbicides. When using ESCORT® XP HERBICIDE on a particular grass for the first time, limit use to a small area. If no injury occurs throughout the season, larger acreage may be treated the following season.

Broadleaf forage species, such as alfalfa and clover, are highly sensitive to ESCORT® XP HERBICIDE and will be severely stunted or injured by ESCORT® XP HERBICIDE.

SPOT TREATMENTS

ESCORT® XP HERBICIDE may be used for use as spot treatment to control noxious and troublesome weeds on pasture, rangeland, and CRP.

Application Information

ESCORT® XP HERBICIDE may be used to control many species of weeds, including noxious weeds, in forage grasses growing on pasture, rangeland, and CRP. Refer to the "Weeds Controlled" section of the package label or supplemental labeling for a listing of susceptible weed species. If the sprayer is calibrated, consult the package label or other supplemental labeling to select the application rate per acre of ESCORT® XP HERBICIDE appropriate for the target weeds. Or mix one gram of ESCORT® XP HERBICIDE per one gallon of water along with a suitable surfactant. Spray to the point of wetting the entire surface of the target weeds, approximately 40 gallons of solution per acre. When applied in this manner there is no grazing restrictions following the use of ESCORT® XP HERBICIDE. Applications may be made at anytime of the year, except when the soil is frozen.

CROP ROTATION

Before using ESCORT® XP HERBICIDE, carefully consider your crop rotation plans and options. For rotational flexibility, do not treat all of your pasture, rangeland, or CRP acres at the same time.

Minimum Rotational Intervals

Minimum rotation intervals* are determined by the rate of breakdown of ESCORT® XP HERBICIDE applied. ESCORT® XP HERBICIDE breakdown in the soil is affected by soil pH, presence of soil microorganisms, soil temperature, and soil moisture. Low soil pH, high soil temperature, and high soil moisture increase ESCORT® XP HERBICIDE breakdown in soil, while high soil pH, low soil temperature, and low soil moisture slow ESCORT® XP HERBICIDE breakdown.

Of these 3 factors, only soil pH remains relatively constant. Soil temperature, and to a greater extent, soil moisture, can vary significantly from year to year and from area to area. For this reason, soil temperatures and soil moisture should be monitored regularly when considering crop rotations.

* The minimum rotation interval represents the period of time from the last application to the anticipated date of the next planting.

Soil pH Limitations

ESCORT® XP HERBICIDE should not be used on soils having a pH above 7.9, as extended soil residual activity could extend crop rotation intervals beyond normal. Under certain conditions, ESCORT® XP HERBICIDE could remain in the soil for 34 months or more, injuring wheat and barley. In addition, other crops planted in high-pH soils can be extremely sensitive to low concentrations of ESCORT® XP HERBICIDE.

Checking Soil pH

Before using ESCORT® XP HERBICIDE, determine the soil pH of the areas of intended use. To obtain a representative pH value for the test area, take several 0" to 4" samples from different areas of the field and analyze them separately. Consult local extension publications for additional information on recommended soil sampling procedures.

BIOASSAY

A field bioassay must be completed before rotating to any crop or grass species/variety not listed in the Rotation Intervals Table, or if the soil pH is not in the specified range, or if the use rate applied is not specified in the table.

To conduct a field bioassay, grow test strips of the crop(s) or grass(es) you plan to grow the following year in fields previously treated with ESCORT® XP HERBICIDE. Crop or grass response to the bioassay will indicate whether or not to rotate to the crop(s) or grass(es) grown in the test strips.

If a field bioassay is planned, check with your local Agricultural dealer or ENVIRONMENTAL SCIENCE U.S., LLC representative for information detailing the field bioassay procedure.

GRAZING/HAYING

When used as directed, there is no grazing or haying restriction for use rates of 1 2/3 ounces per acre and less.

Coveralls, shoes plus socks must be worn if cutting within 4 hours of treatment.

IMPORTANT PRECAUTIONS

- Do not apply more than 1 2/3 ounces of ESCORT® XP HERBICIDE per acre per year on pasture, rangeland, or CRP.
- Grass species or varieties may differ in their response to various herbicides. ENVIRONMENTAL SCIENCE U.S., LLC recommends that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use of ESCORT® XP HERBICIDE to a small area. Components in a grass seed mixture will vary in tolerance to ESCORT® XP HERBICIDE so the final stand may not reflect the seed ratio.
- Under certain conditions such as heavy rainfall, high pH, prolonged cold weather, or wide fluctuations in day/night temperatures prior to or soon after ESCORT® XP HERBICIDE application, temporary discoloration and/or grass injury may occur. ESCORT® XP HERBICIDE should not be applied to grass that is stressed by severe weather conditions, drought, low fertility, water-saturated soils, disease, or insect damage as grass injury may result. Severe winter stress, drought, disease, or insect damage before or following application also may result in grass injury.
- Applications of ESCORT® XP HERBICIDE to pasture, rangeland, and CRP undersown with legumes may cause injury to the legumes. Legumes in a seeding mixture may be severely injured or killed following an application of ESCORT® XP HERBICIDE.
- Applications made to some established grasses may cause temporary stunting, yellowing or seedhead suppression (i.e. fescue, timothy).
- Applications made to newly established grasses less than 2 years from seeding may result in injury or loss.
- Do not apply to forage grasses known to be sensitive to ESCORT® XP HERBICIDE such as ryegrass (Italian and perennial), bahia or Garrison's creeping foxtail.
- Broadleaf forage species, such as alfalfa and clover, are highly sensitive to ESCORT® XP HERBICIDE and will be severely injured or killed.
- The control of weeds in wheel track areas may be reduced if ground applications are made when dry, dusty field conditions exist. The addition of 2,4-D or MCPA should improve weed control under these conditions.

NON-AGRICULTURAL USES

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 700). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

Non-crop industrial weed control and selective weed control in turf (industrial, unimproved only) are not within the scope of the Worker Protection Standard.

NON-CROP SITES

Application Information

ESCORT® XP HERBICIDE is registered for weed control on private, public and military lands as follows: Uncultivated nonagricultural areas (including airports, highway, railroad and utility rights-of-way, sewage disposal areas); uncultivated agricultural areas non-crop producing (including farmyards, fuel storage areas, fence rows, soil bank land, and barrier strips); industrial sites outdoor (including lumberyards, pipeline and tank farms) including grazed areas on these sites. It may also be used for the control of certain noxious and troublesome weeds.

Consult the "Weeds Controlled" and "Brush Species Controlled" tables to determine the appropriate application rate. ESCORT® XP HERBICIDE may be applied in tank mixture with other herbicides labeled for use on non-crop sites. Fully read the labels and follow all directions and restrictions on each label.

Applications may be made by ground or air. Use a sufficient volume of water to ensure thorough coverage of the target vegetation with the application equipment being used.

NATIVE GRASSES

ESCORT® XP HERBICIDE is registered for weed control and suppression in the establishment and maintenance of native grasses. It may be used where blue grama, bluestems (big, little, plains, sand, ww spar) bromegrasses (meadow), buffalograss, green sprangletop, indiangrass, kleingrass, lovegrasses (atherstone, sand, weeping, wilman), orchardgrass, sideoats grama, switchgrass (blackwell), wheatgrass (bluebunch, intermediate, pubescent, Siberian, slender, streamband, tall, thickspike, western), and Russian wildrye are established. It may also be applied over these species in the seedling stage, except for orchardgrass and Russian wildrye.

When used as directed, there are no grazing or hay restrictions for use rates of 1 2/3 ounces per acre or less. At use rates greater than 1 2/3 ounces per acre and up to 3 1/3 ounces per acre, forage grasses may be cut for hay, fodder or green forage and fed to livestock, including lactating animals, 3 days after treatment.

Rotation Intervals for Overseeding and Renovation

Location	Crop or Grass Species	Maximum ESCORT® XP HERBICIDE Rate (oz per A)	Minimum Rotation Interval (months)
AL, AR, FL, GA, KY, LA, MS, NC, OK, SC, TN, TX, VA, WV	Alfalfa, red clover, white clover, sweet clover, bermudagrass, bluegrass, ryegrass, tall fescue	1/10 to 3/10	4
	Wheat (except durum)	1/10 to 3/10	1
	Durum, barley, oat	1/10 to 3/10	10
ALL STATES NOT INCLUDED ABOVE	Red clover, white clover, and sweet clover	1/10 to 2/10	12
	Bermudagrass, bluegrass, ryegrass	1/10 to 2/10	6
	Tall Fescue	1/10 to 2/10	18
	Wheat (except durum)	1/10 to 2/10	1
ALL AREAS WITH SOIL PH OF 7.5 OR LESS	Durum, barley, oat	1/10 to 2/10	10
	Russian wildrye	1/10 to 1/2	1
	Green needlegrass, switchgrass, sheep fescue	1/10 to 1	1
ALL AREAS WITH SOIL PH OF 7.9 OR LESS	Meadow brome, smooth brome, alta fescue, red fescue, meadow foxtail, orchardgrass, Russian wildrye, timothy	1/10 to 1	2
	Alkali sacaton, mountain brome, blue grama, thickspike wheatgrass	1/10 to 1	1
	Sideoats grama, switchgrass	1/10 to 1/2	2
	Western wheatgrass	1/10 to 1	2
	Sideoats grama, switchgrass, big bluestem	1/10 to 1	3

Application Information

Apply ESCORT® XP HERBICIDE at the rate of 1/10 ounce per acre for the control and suppression* of bur buttercup (testiculate), common purslane, common sunflower*, cutleaf eveningprimrose*, flaxweed*, lambsquarters* (common and slimleaf), marehail*, pigweed (redroot and tumble), snow speedwell, tansymustard* and tumble mustard (Jim Hill mustard).

*Suppression is a visual reduction in weed competition (reduced population or vigor) as compared to untreated areas.

Degree of suppression will vary with the size of weed and environmental conditions following treatment.

Application Timing

For established grasses, apply when weeds are in the seedling stage.

For grasses in the seedling stage, apply preplant or preemergence where the soil (seed bed) has been cultivated.

IMPORTANT PRECAUTIONS—NATIVE GRASSES

- Grass species or varieties may differ in their response to various herbicides. If no information is available, limit the initial use of ESCORT® XP HERBICIDE to a small area. Components in a grass seed mixture will vary in tolerance to ESCORT® XP HERBICIDE, so the final stand may not reflect the seed ratio.
- Under certain conditions such as heavy rainfall, high pH, prolonged cold weather, or wide fluctuations in day/night temperatures prior to or soon after ESCORT® XP HERBICIDE application, temporary discoloration and/or grass injury may occur. Injury may result when ESCORT® XP HERBICIDE is applied to grass that is stressed by severe weather conditions, drought, low fertility, water-saturated soils, disease, or insect damage. Severe winter stress, drought, disease, or insect damage before or following application also may result in grass injury.

GRASS REPLANT INTERVALS

Following an application of ESCORT® XP HERBICIDE to non-crop areas, the treated sites may be replanted with various species of grasses at the intervals listed below.

For soils with a pH of 7.5 or less, observe the following replant intervals:

Species	Rate (ounces per acre)	Replant Interval (months)
Brome, Meadow	½-1	2
	1-2	3
Brome, Smooth	½-1	2
	1-2	4
Fescue, Alta	½-1	2
	1-2	4
Fescue, Red	½-1	2
	1-2	4
Fescue, Sheep	½-1	1
	1-2	4
Foxtail, Meadow	½-1	2
	1-2	4
Green Needlegrass	½-2	1
	1-2	4
Orchardgrass	½-1	2
	1-2	4
	2	3
Russian wildrye	½-1	1
	1	2
Switchgrass	½-1	1
	1-2	3
Timothy	½-1	2
	1-2	4
Wheatgrass, Western	½-1	2
	1-2	3

For soils with a pH of 7.5 or greater observe the following replant intervals:

Species	Rate (ounces per acre)	Replant Interval (months)
Alkali Sacaton	½-1	1
	1-2	3
Bluestem, Big	½-2	3
	1-2	1
Brome, Mountain	½-1	1
	1-2	2
Grama, Blue	½-2	1
	½	2
Grama, Sideoats	½	2
	>½	>3
Switchgrass	½	2
	>½	>3
Wheatgrass, Thickspike	½-2	1
Wheatgrass, Western	1-2	2
	½-1	3

The specified intervals are for applications made in the Spring to early Summer. Because ESCORT® XP HERBICIDE degradation is slowed by cold or frozen soils, applications made in the late Summer or Fall should consider the intervals as beginning in the Spring following treatment.

Testing has indicated that there is considerable variation in response among the species of grasses when seeded into areas treated with ESCORT® XP HERBICIDE. If species other than those listed above are to be planted into areas treated with ESCORT® XP HERBICIDE, a field bioassay must be performed, or previous experience may be used, to determine the feasibility of replanting treated sites.

ADDITIONAL GRASS INFORMATION

APPLICATION INFORMATION FOR GRASS ESTABLISHMENT

ESCORT® XP HERBICIDE may be used for the control or suppression of broadleaf weeds to aid in the establishment of the following perennial native or improved grasses:

Blue Gramma	Buffalograss	Orchardgrass	Wheatgrasses –	Wildrye grass –
Bluestems –	Green sprangletop	Sideoats gramma	bluebunch	Russian
big	Kleingrass	Switchgrass –	crested	
little	Lovegrasses –	Blackwell	intermediate	
plains	Atherstone		pubescent	
sand	sand		Siberian	
WW spar	weeping		slender	
	Wilman		steambank	
			tall	
			thickspike	
			western	

Maximize potential for grass establishment by consulting with the Natural Resource and Conservation Service of other government agencies or local experts concerning planting techniques and other cultural practices.

Performance from ESCORT® XP HERBICIDE may not always be satisfactory due to the inability of newly planted grass stands to sufficiently compete with weeds and the severity of weed pressure in new grass stands.

An additional herbicide application or mowing may be needed.

Use Rates and Application Timing for Grass Establishment Preplant (prior to planting) or Preemergence (after planting but before grass emergence)

Do not use more than 1/10 ounce per acre of ESCORT® XP HERBICIDE for grass establishment.

Apply ESCORT® XP HERBICIDE at 1/10 ounce per acre on all labeled grasses except orchardgrass and Russian wildrye grass. Do not apply ESCORT® XP HERBICIDE preplant or preemergence to orchardgrass and Russian wildrye grass as severe crop injury may result.

Early postemergence to new plantings

Apply ESCORT® XP HERBICIDE at 1/10 ounce per acre, plus a non-ionic surfactant at the rate of 2 to 4 pints per 100 gallons of spray solution on all labeled grasses anytime after grass emergence.

Do not use a spray adjuvant other than non-ionic surfactant.

Because grass species differ in time of emergence, apply only after the majority of grasses are in the 3 to 4 leaf stage.

Postemergence to stands with 1 – 5 leaf grasses planted the previous season

Apply ESCORT® XP HERBICIDE at 1/10 ounce per acre plus a non-ionic surfactant at the rate of 2 to 4 pints per 100 gallons of spray solution, on all labeled grasses when the majority of the grasses have one or more leaves.

Do not use a spray adjuvant other than non-ionic surfactant.

APPLICATION INFORMATION FOR ESTABLISHED GRASSES

Use Rates for Established Grasses

Apply up to 1 ounce ESCORT® XP HERBICIDE per acre as a broadcast application to established grasses. For spot applications, use 1 ounce per 100 gallons of water. Do not apply more than 1 2/3 ounces of ESCORT® XP HERBICIDE per acre per year.

Refer to the Weeds Controlled section of this label for a listing of the weeds controlled by ESCORT® XP HERBICIDE and the appropriate use rate to obtain control.

Application Timing – Established Grasses

ESCORT® XP HERBICIDE may be applied to established native grasses such as bluestems and grama, and on other established grasses such as bermudagrass, bluegrass, orchardgrass, bromegrass, fescue and timothy that were planted the previous growing season (or earlier) and are fully tillered, unless otherwise directed on this label. Specific application timing information on several of these grass species follows:

Minimum time from Grass establishment

ESCORT® XP HERBICIDE application

Grass	
Bermudagrass	2 months
Bluegrass, bromegrass, Orchardgrass	6 months
Timothy	12 months
Fescue	24 months

Fescue and Timothy Precautions

When used on fescue and timothy grasses, ESCORT® XP HERBICIDE may cause reduced first cutting yields due to temporary stunting, leaf yellowing, or seed head suppression. To help minimize these symptoms, follow the information below:

- Use the lowest labeled rate for the target weeds.
- Tank mix 2,4-D with ESCORT® XP HERBICIDE applications.
- Apply ESCORT® XP HERBICIDE at no more than 4/10 ounce per acre.
- Make applications when the grasses are 5 to 6 inches tall in late summer or fall.
- Use only a non-ionic surfactant at 1/2 pint per 100 gallons of spray solution.
- When liquid nitrogen is the spray carrier, do not include the surfactant.

Other Grasses:

Application of ESCORT® XP HERBICIDE to Pensacola bahiagrass, ryegrass (Italian or perennial) and Garrison's creeping foxtail may cause severe injury to and/or loss of forage.

Varieties and species of forage grasses differ in their tolerance to herbicides. When using ESCORT® XP HERBICIDE on a particular grass for the first time, limit use to a small area. If no injury occurs throughout the season, larger acreage may be treated the following season.

Broadleaf forage species, such as alfalfa and clover, are highly sensitive to ESCORT® XP HERBICIDE and will be severely stunted or injured by ESCORT® XP HERBICIDE.

CROP ROTATION

Before using ESCORT® XP HERBICIDE, carefully consider your crop rotation plans and options.

Minimum Rotational Intervals

Minimum rotation intervals* are determined by the rate of breakdown of ESCORT® XP HERBICIDE applied. ESCORT® XP HERBICIDE breakdown in the soil is affected by soil pH, presence of soil microorganisms, soil temperature, and soil moisture. Low soil pH, high soil temperature, and high soil moisture increase ESCORT® XP HERBICIDE breakdown in soil, while high soil pH, low soil temperature, and low soil moisture slow ESCORT® XP HERBICIDE breakdown.

Of these 3 factors, only soil pH remains relatively constant. Soil temperature, and to a greater extent, soil moisture, can vary significantly from year to year and from area to area. For this reason, monitor soil temperature and soil moisture on a regular basis when considering any crop rotations.

* The minimum rotation interval represents the period of time from the last application to the anticipated date of the next planting.

Soil pH Limitations

ESCORT® XP HERBICIDE must not be used on soils having a pH above 7.9, as extended soil residual activity could extend crop rotation intervals beyond normal. Under certain conditions, ESCORT® XP HERBICIDE could remain in the soil for 34 months or more, injuring wheat and barley. In addition, other crops planted in high-pH soils can be extremely sensitive to low concentrations of ESCORT® XP HERBICIDE.

Checking Soil pH

Before using ESCORT® XP HERBICIDE, determine the soil pH of the areas of intended use. To obtain a representative pH value for the test area, take several 0" to 4" samples from different areas of the field and analyze them separately. Consult local extension publications for additional information on recommended soil sampling procedures.

BIOASSAY

A field bioassay must be completed before rotating to any crop or grass species/variety not listed in the Rotation Intervals Table, or if the soil pH is not in the specified range, or if the use rate applied is not specified in the table.

To conduct a field bioassay, grow test strips of the crop(s) or grass(es) you plan to grow the following year in fields previously treated with ESCORT® XP HERBICIDE. Crop or grass response to the bioassay will indicate whether or not to rotate to the crop(s) or grass(es) grown in the test strips.

If a field bioassay is planned, check with your local Agricultural dealer or ENVIRONMENTAL SCIENCE U.S., LLC representative for information detailing the field bioassay procedure.

IMPORTANT PRECAUTIONS

- Grass species or varieties may differ in their response to various herbicides. If no information is available, limit the initial use of ESCORT® XP HERBICIDE to a small area.
- Components in a grass seed mixture will vary in tolerance to ESCORT® XP HERBICIDE so the final stand may not reflect the seed ratio.
- Under certain conditions, such as heavy rainfall, high pH, prolonged cold weather, or wide fluctuations in day/night temperatures, prior to or soon after ESCORT® XP HERBICIDE application, temporary discoloration and/or grass injury may occur. ESCORT® XP HERBICIDE applied to grass that is stressed by severe weather conditions, drought, low fertility, water-saturated soils, disease, or insect damage can result in grass injury. Severe winter stress, drought, disease, or insect damage before or following application also may result in grass injury.
- Applications of ESCORT® XP HERBICIDE to lands underdown with legumes may cause injury to the legumes. Legumes in a seeding mixture may be severely injured or killed following an application of ESCORT® XP HERBICIDE.
- The control of weeds in wheel track areas may be reduced if ground applications are made when dry, dusty field conditions exist. The addition of 2,4-D or MCPA may improve weed control under these conditions.

WEEDS CONTROLLED**1/3 to 1/2 ounce per acre**

Annual sowthistle	Common groundsel	Goldenrod	Smallseed falseflax
Aster	Common purslane	Lambsquarters	Smooth pigweed
Bahiagrass	Common yarrow	Marestail/horseweed****	Sweet clover
Beebalm	Conical catchfly	Maximillion sunflower	Tansymustard
Bittercress	Corn cockle	Miners lettuce	Teacle mustard
Bitter sneezeweed	Cow cockle	Pennsylvania smartweed	Tumble mustard
Blackeyed-susan	Crown vetch	Plains coreopsis	Wild carrot
Blue mustard	Dandelion	Plantain	Wild garlic
Bur buttercup	Dogfennel	Redroot pigweed	Wild lettuce
Chicory	False chamomile	Redstem filaree	Wild mustard
Clover	Fiddleneck tarweed	Rough fleabane	Wooly croton
Cocklebur	Field pennycress	Shepherd's purse	Wood sorrel
Common chickweed	Flxweed	Silky crazyweed (locoweed)	Yankweed

1/2 to 1 ounce per acre

Blackberry	Common sunflower	Henbit	Rosering gaillardia
Black henbane	Curly dock	Honeysuckle	Seaside arrowgrass
Broom snakeweed*	Dewberry	Multiflora rose and other wild roses	Sericea lespedeza
Buckhorn plantain	Dyer's woad	Musk thistle***	Tansy ragwort
Bull thistle	Garlic mustard	Oxeye daisy	Teasel
Common crupina	Gorse	Plumless thistle	Wild caraway
	Halogeton	Prostrate knotweed	

1 to 2 ounces per acre

Common mullein	Lupine	Purple scabious	Sulphur cinquefoil
Common tansy	Old world climbing fern	Scotch thistle	Western salsify
Field bindweed**	(<i>Lygodium</i>)	Scouringrush	Whitetop (hoary cress)
Greasewood	Perennial pepperweed	Salsify	Wild Iris
Gumweed	Poison hemlock	Snowberry	
Houndstongue	Purple loosestrife	St. Johnswort	

1 1/2 to 2 ounces per acre

Canada thistle**	Duncecap larkspur	Tall larkspur	Yellow toadflax**
Dalmation toadflax**	Russian Knapweed**	Wild parsnip	

2 ounces per acre

Onionweed

3 to 4 ounces per acre

Kudzu

* Apply fall through spring.

** Suppression, which is a visual reduction in weed competition (reduced population or vigor) as compared to untreated areas. Apply as a full coverage spray for best performance.

*** Certain biotypes of musk thistle are more sensitive to ESCORT® XP HERBICIDE and may be controlled with rates of 1/4 to 1/2 ounce per acre. Treatments of ESCORT® XP HERBICIDE may be applied from rosette through bloom stages of development.

**** Certain biotypes of marestail/horsetail are less sensitive to ESCORT® XP HERBICIDE and may be controlled by tank mixes with herbicides with a different mode of action.

Problem Weed Control

For broader spectrum control and for use on certain biotypes of broadleaf weeds which may be resistant to ESCORT® XP HERBICIDE and herbicides with the same mode of action, the following tank mixes may be used.

Dicamba + 2,4-D

Weed	Rate of ESCORT® XP HERBICIDE	Rate of dicamba (fluid ounces/acre)	Rate of 2,4-D (fluid ounces/acre)
Kochia control	1/2	8	16
Spotted knapweed control	1/2	8	16
Rush skeletonweed suppression	1	8	16

INDUSTRIAL TURFGRASS UNIMPROVED ONLY**Application Information**

ESCORT® XP HERBICIDE is registered for selective weed control in unimproved industrial turfgrass where certain grasses are well established and desired as ground cover. ESCORT® XP HERBICIDE may also be used for the control of certain noxious and troublesome weeds in turfgrass.

In addition to conventional spray equipment, ESCORT® XP HERBICIDE may also be applied with invert emulsion equipment. When using an invert emulsion, mix the prescribed rate of ESCORT® XP HERBICIDE in the water phase.

Consult the "Weeds Controlled" table to determine which weeds will be controlled by the following application rates:

Turfgrass Type	Rate of ESCORT® XP HERBICIDE (ounces/acre)
Fescue and Bluegrass	¼ to ½
Crested Wheatgrass and Smooth Brome	¼ to 1
Bermudagrass	¼ to 2

Application Timing

Applications may be made at anytime of the year except when the soil is frozen.

When a spring application is made on fescue or bluegrass, a second application may be made during the summer after full seedhead maturation.

Growth Suppression and Seedhead Inhibition (Chemical Mowing)**Application Information**

ESCORT® XP HERBICIDE may be used for growth suppression and seedhead inhibition in well established fescue and bluegrass turfgrass at the use rate of 1/4 to 1/2 ounce per acre.

Tank Mix Combination

ESCORT® XP HERBICIDE may be tank mixed with "Embank" for improved performance in the regulation of growth and seedhead suppression. Tank mix 1/4 to 1/2 ounce of ESCORT® XP HERBICIDE with 1/8 to 1/4 pint of "Embank".

Application Timing

Application may be made after at least 2 to 3 inches of new growth has emerged until the appearance of the seed stalk.

IMPORTANT PRECAUTIONS**—INDUSTRIAL TURFGRASS ONLY**

- An application of ESCORT® XP HERBICIDE may cause temporary discoloration (chlorosis) or stunting of the turfgrasses. Use the lower specified rates for minimum discoloration or stunting.
- With fescue and bluegrass, sequential applications made during the same or consecutive growth periods (i.e. spring and fall) may result in excessive injury to turfgrass.
- Excessive injury may result when ESCORT® XP HERBICIDE is applied to turfgrass that is under stress from drought, insects, disease, cold temperatures (winter injury) or poor fertility.
- ESCORT® XP HERBICIDE is not recommended for use on bahiagrass.

BRUSH CONTROL**Application Information**

ESCORT® XP HERBICIDE is registered for the control of undesirable brush growing in non-crop areas including grazed areas on these sites. Applications may be made by air, high volume ground application, low volume ground application and ultra-low volume ground application. Except as noted for multiflora rose, ESCORT® XP HERBICIDE must be applied as a spray to the foliage.

The application volume required will vary with the height and density of the brush and the application equipment used. Generally, aerial applications will require 15 to 25 gallons of water per acre; high volume ground application will require 100 to 400 gallons of water per acre; low volume ground application will require 20 to 50 gallons of water per acre; and ultra-low volume ground application will require 10 to 20 gallons of water per acre.

Regardless of the application volume and equipment used, thorough coverage of the foliage, particularly the terminal growing points, is necessary to optimize results.

BRUSH SPECIES CONTROLLED

Species	High Volume Rate (ounces/100 gallon)	Broadcast Rate (ounces/acre)
Ash	1-2	1-3
Aspen	1-2	1-3
Black locust	1-2	1-3
Blackberry	1-2	1-3
Camelthorn	1-2	1-3
Cherry	1-2	1-3
Cottonwood	1-2	2-3
Eastern red cedar	1-2	2-3
Elder	1-2	2-3
Elm	1-2	1-3
Firs	3	1-2
Hawthorn	1-2	1-3
Honeysuckle	1-2	1/2-1
Mulberry	1-2	2-3
Multiflora rose	1-2	1-3
Muscadine (wild grape)	1-2	2-3
Oaks	1-2	1-3
Ocean spray (Holodiscus)	1-2	2-3
Osage orange	1-2	2-3
Red maple	1-2	1-3
Salmonberry	1/2-1	1-3
Snowberry	1/2-1	1-3
Spruce (black and white)	3	2-3
Thimbleberry	1/2-1	1-3
Tree of heaven (Ailanthus)	1-2	1-2
Wild roses	1/2-1	1-3
Willow	1/2-1	1-3
Yellow poplar	1/2-1	1-3

For low volume and ultra-low volume ground applications, mix 4 to 8 ounces of ESCORT® XP HERBICIDE per 100 gallons of spray solution.

Application Timing

Make a foliar application of the specified rate of ESCORT® XP HERBICIDE during the period from full leaf expansion in the spring until the development of full fall coloration on deciduous species to be controlled. Coniferous species may be treated at anytime during the growing season.

Spot Treatment

ESCORT® XP HERBICIDE may be used for the control of many species of weeds including noxious/invasive weeds in certain established grasses growing on non-crop areas.

Refer to the "Weeds Controlled" section for a listing of susceptible weed species and the application rate per acre per the target weed.

Or, mix one gram of ESCORT® XP HERBICIDE per one gallon of water along with a surfactant. Spray to the point of wetting the entire surface of the target weeds, approximately 40 gallons of solution per acre.

Tank Mix Combinations—

ESCORT® XP HERBICIDE may be tank mixed with any product labeled for non-crop brush control at the application rates specified on the companion product's label for the pests specified on the product's companion label. Read and follow the label instructions of both products when tank mixing. Follow the most restrictive limitations of any of the product labels being tank mixed.

Low Rate Applications

Imazapyr (2 pound active per gallon)

Combine 1 to 2 ounces of ESCORT® XP HERBICIDE with 1 to 4 pints of imazapyr herbicide per acre and apply as a broadcast spray. For aerial applications use a minimum of 15 gallons per acre spray volume. In addition to species listed above controlled by ESCORT® XP HERBICIDE, this combination controls black gum, hophornbeam, sassafras, sweetgum, Vaccinium species, dogwood, myrtle dahoon, hickories, and persimmon.

Picloram* (2 pound active per gallon) + Imazapyr (2 pound active per gallon)

Combine 1 to 1 1/2 ounce of ESCORT® XP HERBICIDE with 2 to 8 fluid ounces of imazapyr and 1 to 2 pints of picloram per 100 gallons of water. Apply as a high volume spray. This tank mix controls cherry, elms, box elder, maples, hackberry, redbud, ash, oaks (including shingle oak), black locust, and sassafras.

*Picloram is a restricted use pesticide.

Spotgun Basal Soil Treatment

For control of multiflora rose, prepare a spray suspension of ESCORT® XP HERBICIDE by mixing 1 ounce per gallon of water. Mix vigorously until the ESCORT® XP HERBICIDE is dispersed and agitate periodically while applying the spray suspension.

Apply the spray preparation with an exact delivery handgun applicator. Apply at the rate of 4 milliliters for each 2 feet of rose canopy diameter. Direct the treatment to the soil within 2 feet of the stem union. When treating large plants and more than one delivery is required, make applications on opposite sides of the plant.

For best results, make applications from early spring to summer.

IMPORTANT PRECAUTIONS

—NON-CROP BRUSH ONLY

• When using tank mixtures of ESCORT® XP HERBICIDE with companion herbicides, read and follow all use instructions, application rates, warnings, and precautions appearing on the labels. Follow the most restrictive label instructions for each of the herbicides used.

SPRAY EQUIPMENT

Low rates of ESCORT® XP HERBICIDE can kill or severely injure most crops. Following an ESCORT® XP HERBICIDE application, the use of spray equipment to apply other pesticides to crops on which ESCORT® XP HERBICIDE is not registered may result in their damage. The most effective way to reduce this crop damage potential is to use dedicated mixing and application equipment.

MIXING INSTRUCTIONS

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of ESCORT® XP HERBICIDE.
3. Continue agitation until the ESCORT® XP HERBICIDE is fully dispersed, at least 5 minutes.
4. Once the ESCORT® XP HERBICIDE is fully dispersed, maintain agitation and continue filling tank with water. ESCORT® XP HERBICIDE must be thoroughly mixed with water before adding any other material.
5. As the tank is filling, add tank mix partners (if desired) then add the necessary volume of nonionic surfactant. Always add surfactant last.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. ESCORT® XP HERBICIDE spray preparations are stable if they are pH neutral or alkaline and stored at or below 100° F.
8. If ESCORT® XP HERBICIDE and a tank mix partner are to be applied in multiple loads, pre-slurry the ESCORT® XP HERBICIDE in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the ESCORT® XP HERBICIDE.

PRODUCT PRECAUTIONS

- When used as directed, there is no grazing or haying restriction for use rates of 1 2/3 ounces per acre or less. At use rates greater than 1 2/3 ounces per acre and up to 3 1/3 ounces per acre, forage grasses may be cut for hay, fodder or green forage and fed to livestock, including lactating animals, 3 days after treatment.
- Injury to or loss of desirable trees or other plants may result if spray equipment is drained or flushed on or near these trees or plants, or on areas where their roots may extend, or in locations where the product may be washed or moved into contact with their roots.
- Treatment of powdery, dry soil or light, sandy soil when there is little likelihood of rainfall soon after treatment may result in off target movement and possible damage to susceptible crops when soil particles are moved by wind or water. Injury to crops may result if treated soil is washed, blown, or moved onto land used to produce crops. Exposure to ESCORT® XP HERBICIDE may injure or kill most crops. Injury may be more severe when the crops are irrigated. Do not apply ESCORT® XP HERBICIDE when these conditions are identified and powdery, dry soil or light or sandy soils are known to be prevalent in the area being treated.
- Applications made where runoff water flows onto agricultural land may injure crops. Applications made during periods of intense rainfall, to soils saturated with water, to surfaces paved with materials such as asphalt or concrete, or to soils through which rainfall will not readily penetrate may result in runoff and movement of ESCORT® XP HERBICIDE.
- Do not treat frozen or snow covered soil.
- Leave treated soil undisturbed to reduce the potential for ESCORT® XP HERBICIDE movement by soil erosion due to wind or water.

PRODUCT RESTRICTIONS

- Do not use on lawns, walks, driveways, tennis courts, or similar areas.
- Do not apply through any type of irrigation system.
- Do not use this product in the following counties of Colorado: Saguache, Rio Grande, Alamosa, Costilla and Conejos.
- Do not use this product in California

SPRAYER CLEANUP

Spray equipment must be cleaned before ESCORT® XP HERBICIDE is sprayed. Follow the cleanup procedures specified on the labels of previously applied products. If no directions are provided, follow the six steps outlined below.

When multiple loads of ESCORT® XP HERBICIDE are applied, it is recommended that at the end of each day of spraying, the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits that can accumulate in the application equipment.

1. Drain tank; thoroughly rinse spray tanks, boom, and hoses with clean water. Loosen and physically remove any visible deposits.
2. Fill the tank with clean water and 1 gallon of ammonia (contains 3% active minimum) for every 100 gallons of water. Flush the hoses, boom, and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 minutes. Flush the hoses, boom, and nozzles again with the cleaning solution, and then drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.

- Repeat step 2.
- Rinse the tank, boom, and hoses with clean water.
- Dispose of the rinsate on a labeled site or at an approved waste disposal facility. If a commercial cleaner is used follow the commercial cleaner directions for rinsate disposal.

Notes:

- Mixing chlorine bleach with ammonia can cause dangerous gases to form. Clean spray equipment outdoors.
- Use steam cleaning or other commercial cleaners to facilitate the removal of any caked pesticide deposits.
- When ESCORT® XP HERBICIDE is tank mixed with other pesticides, all cleanout procedures for each product must be examined and the most rigorous procedure must be followed.
- In addition to this cleanout procedure, all pre-cleanout guidelines on subsequently applied products must be followed as per the individual product labels.

SPRAY DRIFT MANAGEMENT

Aerial Applications:

- Do not release spray at a height greater than 10 ft above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Ground Boom Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or target vegetation unless making an industrial turf, pasture and rangeland applications, in which case applicators may apply with a nozzle height no more than 4 feet above the crop or target vegetation.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Boom-less Ground Applications:

- Applicators are required to use a Medium or coarser droplet size (ASABE S572.1) for all applications.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

Boom-less Ground Applications:

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles - Follow nozzle manufacturer recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift. When applying aerially to crops, do not release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, that it is configured properly, and that drift potential has been minimized.

Note: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Read the specific crop use and application equipment instructions to determine if an air assisted field crop sprayer can be used.

SENSITIVE AREAS

Making applications when there is a sustained wind moving away from adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is an effective way to minimize the effect of spray drift.

DRIFT CONTROL ADDITIVES

Using product compatible drift control additives can reduce drift potential. When a drift control additive is used, read and carefully observe cautionary statements and all other information on the additive's label. If using an additive that increases viscosity, ensure that the nozzles and other application equipment will function properly with a viscous spray solution. Preferred drift control additives have been certified by the Chemical Producers and Distributors Association (CPDA).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store product in original container only. Store in a cool, dry place.

Pesticide Disposal: Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling:

Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Nonrefillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers (Capacity Greater Than 50 Pounds): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down): Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom, and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume.

(continued)

STORAGE AND DISPOSAL *(continued)*

If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Paper or Plastic Bags, Fiber Sacks including Flexible Intermediate Bulk Containers (FIBC) or Fiber Drums With Liners: Nonrefillable container. Do not reuse or refill this container. Completely empty paper or plastic bag, fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

Refillable Fiber Drums With Liners: Refillable container (fiber drum only). Refilling Fiber Drum: Refill this fiber drum with ESCORT® XP HERBICIDE containing metsulfuron methyl only. Do not reuse this fiber drum for any other purpose. Cleaning before refilling is the responsibility of the refiller. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Disposing of Fiber Drum and/or Liner: Do not reuse this fiber drum for any other purpose other than refilling (see preceding). Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container. Offer the liner for recycling if available or dispose of liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner. To clean the fiber drum before final disposal, completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

All Other Refillable Containers: Refillable container. Refilling Container: Refill this container with ESCORT® XP HERBICIDE containing metsulfuron methyl only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, do not use the container, contact ENVIRONMENTAL SCIENCE U.S., LLC at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, do not reuse or transport container, contact ENVIRONMENTAL SCIENCE U.S., LLC at the number below for instructions. Disposing of Container: Do not reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom, and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour, or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Outer Foil Pouches of Water Soluble Packets (WSP): Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or, dispose of the empty outer foil pouch in the trash as long as WSP is unbroken. If the outer pouch contacts the formulated product in any way, the pouch must be triple rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer pouch as described previously. Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact ENVIRONMENTAL SCIENCE U.S., LLC at 1-800-424-9300, day or night.

"Embank" is a registered trademark of PBI Gordon Corporation.

Envu and the Envu logo are trademarks and Escort® is a registered trademark owned by Environmental Science U.S., LLC or one of its affiliates.

Velpar is a registered trademark of E.I Dupont de Nemours and company used under license by Environmental Science U.S., LLC.

CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, plant injury, other property damage, as well as other unintended consequences may result because of factors beyond the control of Environmental Science U.S., LLC. Those factors include, but are not limited to, weather conditions, presence of other materials or the manner of use or application. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ENVIRONMENTAL SCIENCE U.S., LLC MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Environmental Science U.S., LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ENVIRONMENTAL SCIENCE U.S., LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT ENVIRONMENTAL SCIENCE U.S., LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.

For product information call: 1-800-331-2867



ESCORT[®]

XP

Herbicide

Dry Flowable

Active Ingredient

Metsulfuron methyl
Methyl 2-[[[4-methoxy-6-methyl-1,3,5-triazin-
2-yl)amino]carbonyl]amino]sulfonyl]benzoate..... 60%

Other Ingredients 40%

TOTAL

EPA Reg. No. 101563-167

By Weight

100%

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

See Panel for First Aid Instructions for Complete Precautionary Statements and Directions for Use.

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! Causes eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

This herbicide is injurious to plants at extremely low concentrations. Nontarget plants may be adversely effected from drift and run-off.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store product in original container only. Store in a cool, dry place.

Pesticide Disposal: Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling:

Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Nonrefillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds):

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact ENVIRONMENTAL SCIENCE U.S., LLC at 1-800-424-9300, day or night.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact ENVIRONMENTAL SCIENCE U.S., LLC at 1-800-424-9300, day or night.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact ENVIRONMENTAL SCIENCE U.S., LLC at 1-800-424-9300, day or night.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact ENVIRONMENTAL SCIENCE U.S., LLC at 1-800-424-9300, day or night.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact ENVIRONMENTAL SCIENCE U.S., LLC at 1-800-424-9300, day or night.

Envu and the Envu logo are trademarks and Escort[®] is a registered trademark owned by Environmental Science U.S., LLC or one of its affiliates.

PRODUCED FOR

Environmental Science U.S., LLC
5000 CentreGreen Way, Suite 400
Cary, NC 27513

85798669

85796941D 221216AV1

Nonrefillable Container

Net Weight:

1 Pound

envu[™]

