

◀ PULL HERE TO OPEN

GROUP 29 HERBICIDE



Esplanade™ 200 SC

Preemergent Herbicide for the Control of Annual Grasses and Broadleaf Weeds in Non-Residential Non-Crop Areas, Railroad and Rail Yards, Managed Roadsides, Fence Rows, Utilities, Hardscapes, Industrial, Municipal, and Government Sites

ACTIVE INGREDIENT:

Indaziflam (CAS No: 730979-19-8) 19.05%

OTHER INGREDIENTS: 80.95%

Total: 100.00%

Contains 1.67 pounds of indaziflam per gallon.

EPA Reg. No. 432-1516 EPA Est. No. 000264-DEU-001

KEEP OUT OF REACH OF CHILDREN CAUTION

For MEDICAL And TRANSPORTATION Emergencies

ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-800-331-2867

Refer to attached leaflet for complete Directions for Use and Precautionary Statements.

BACKED
by **BAYER™**

Net Contents

2.5 Gallons (9.46 L)

80238924

80522975A

110303A



Produced for:

Bayer Environmental Science

A Division of Bayer CropScience LP

2 T. W. Alexander Drive

Research Triangle Park, NC 27709

Product of Germany

Esplanade™ 200 SC

Preemergent Herbicide for the Control of Annual Grasses and Broadleaf Weeds in Non-Residential Non-Crop Areas, Railroad and Rail Yards, Managed Roadsides, Fence Rows, Utilities, Hardscapes, Industrial, Municipal, and Government Sites

ACTIVE INGREDIENT:

Indaziflam (CAS No: 730979-19-8).....19.05%

OTHER INGREDIENTS:.....80.95%

Total.....100.00%

EPA Reg No. 432-1516

EPA Est. No. 000264-DEU-001

Contains 1.67 pounds of indaziflam per gallon

KEEP OUT OF REACH OF CHILDREN CAUTION

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-800-334-9745

FIRST AID

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> • Hold eyes open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
<p>For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577 Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p>	

PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, absorbed through the skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This Product is toxic to fish, aquatic invertebrates, and plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean water mark. Do not contaminate water when disposing of rinseate or washwater. This product may impact water through spray drift or runoff. Follow directions for use to avoid spray drift and runoff. A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential of this product entering water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Ground Water Advisory: This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This pesticide may impact water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Protect Esplanade 200 SC Herbicide from freezing temperatures.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are acutely hazardous.

Improper disposal of excess pesticide, spray mixture, or rinseate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinseate into application equipment or a mix tank or store rinseate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available or reconditioning, if appropriate. Then puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

Refer to attached leaflet for complete Directions for Use and Precautionary Statements.

**Net Contents
2.5 Gallons (9.46 L)
80238924
80522975A 110303A**



Produced for:

Bayer Environmental Science

A Division of Bayer CropScience LP

2 T. W. Alexander Drive

Research Triangle Park, NC 27709

Product of Germany

FIRST AID

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> • Hold eyes open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
<p>For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577 Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, absorbed through the skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators and other handlers must wear:

- long-sleeved shirt and long pants
- shoes plus socks
- chemical resistant gloves made of any waterproof material such as natural rubber > 14 mils

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS:

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d) (4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

This Product is toxic to fish, aquatic invertebrates, and plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean water mark. Do not contaminate water when disposing of rinsate or washwater. This product may impact water through spray drift or runoff. Follow directions for use to avoid spray drift and runoff. A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential of this product entering water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Ground Water Advisory: This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This pesticide may impact water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read the entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

SHAKE CONTAINER WELL BEFORE USING.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes plus socks
- Chemical resistant gloves made of any waterproof material

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Protect Esplanade 200 SC Herbicide from freezing temperatures.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are acutely hazardous.

Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as

Wind

Avoid making applications when spray particles may be carried by air currents to areas where sensitive crops and plants are growing. Many factors influence spray drift potential including droplet size, equipment type and local terrain. Drift potential increases if wind is in excess of 10 mph, gusty, or below 2 mph (due to inversion potential). Always make applications when there is some air movement to determine the direction and distance of possible spray drift. The applicator should be familiar with local conditions and how it may influence spray drift.

Temperature Inversion

A surface temperature inversion (i.e., increasing temperature with increasing altitude) greatly increases the potential for drift. Avoid application when conditions are favorable to inversion. Presence of ground fog is a good indicator of a surface temperature inversion.

Controlling Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that still provide sufficient coverage and control. Uniform spray coverage is important to maximize weed control. Applying larger droplets will reduce drift potential, but will not prevent drift if applications are made improperly or under unfavorable environmental conditions such as wind speed, temperature and humidity, and temperature inversion situations.

Spray volume, pressure, and nozzle selection are all important for reducing drift. Select a high flow rate nozzle to apply the highest practical spray volume. High flow rate nozzles produce larger droplets. Use lower spray pressures within the recommended range for the nozzle. If a higher flow rate is needed, increase the nozzle size instead of increasing pressure. Lower spray pressures produce larger droplets. Also, consider using low-drift nozzles.

Set the boom and make applications at the lowest height that safely permits uniform coverage of the soil and minimizes droplet evaporation. Avoid application if wind conditions are gusty. Local terrain may influence wind patterns. The applicator should be familiar with local conditions and understand how they may impact spray drift.

Drift Control Additive

Drift control additive may also be used with most spray equipment to reduce the potential for drift. When using a drift control additive, read and follow all directions on the additive label.

Shielded Sprayers

Shielding the boom or individual nozzles may also reduce the potential for drift. However, it is the responsibility of the applicator to verify that the shield does not interfere with uniform spray coverage.

Vegetation Management Information

Timings, Use Rates, and Maximum Seasonal Rate for Esplanade 200 SC

Apply Esplanade 200 SC prior to weed seed germination. Esplanade 200 SC does not generally control weeds that have emerged. For maximum weed control, the herbicide needs to reach the soil surface and be activated by rainfall or adequate soil moisture. Apply Esplanade 200 SC in the spring for control of spring and summer germinating weeds and apply in the fall for control of winter weeds.

The desired rate of Esplanade 200 SC depends on the residual weed activity required and restrictions on the maximum amount of Esplanade 200 SC per season. Esplanade 200 SC may be applied at 3.5- 7 fl oz per acre. Do not exceed 7 fl oz of Esplanade 200 SC for a single application. Applications of Esplanade 200 SC must not exceed the maximum label rate (10 fl oz per acre) in a 12 month period after the previous application.

Factors including soil type, rainfall, and the amount of vegetation at the time of treatment may affect weed control. Lower rates of Esplanade 200 SC may be effective for sandy soils, whereas organic soils may require higher rates. If the herbicide is not activated by rainfall prior to weed germination, control may be reduced.

For late fall applications, apply Esplanade 200 SC prior to when the ground freezes.

Tank Mix Combinations

Tank-mix combinations of Esplanade 200 SC plus a non-selective herbicide such as Finale Herbicide or glyphosate will control existing undesirable vegetation in dormant warm season grasses. Applied as a broadcast spray, Esplanade 200 SC plus a non-selective herbicide such as Finale Herbicide or glyphosate will provide pre and post-emergent control of susceptible species listed on the respective labels of the herbicides in the tank mixture.

Esplanade 200 SC may be tank mixed with following herbicide active ingredients but not limited to: [2,4-D, aminopyralid, bromacil, dicamba, flumioxazin, fosamine, glufosinate ammonium (Finale Herbicide), glyphosate, hexazinone, metsulfuron, picloram, simazine, sulfometuron, and triclopyr]

Follow all use restrictions on this label and for all tank mix partners and use the most restrictive use pattern for the labels of all products in a tank mixture.

Apply mixtures so that the spray solution covers the soil surface in a uniform manner. If uniform coverage is not achieved, preemergent activity will be inconsistent.

Resistance Management Guidelines

Continual use of herbicides with a single mode of action encourages the development of resistant weeds. Esplanade 200 SC Herbicide is a Group 29 Herbicide that contains the active ingredient indaziflam. Esplanade 200 SC Herbicide may be used in programs with other preemergence herbicides with different modes of action. No known resistance to Esplanade 200 SC Herbicide exists, and there are no known instances of cross resistance between this product and other classes of herbicides, or modes of action. Performance of this product is not affected by the presence of biotypes resistant to glyphosate, triazines, ALS-inhibiting, growth regulator, or other herbicide modes of action. When resistance of a specific weed is confirmed, rotation of Esplanade 200 SC Herbicide in one season followed by a preemergent herbicide with another mode of action in the subsequent season, for example, will reduce existing populations and minimize further development of resistant weeds. Contact a Bayer Environmental Sciences representative for the latest information on resistance management guidelines for this product.

Weeds Controlled or Suppressed by Esplanade 200SC

Broadleaf Weeds Controlled

American black nightshade Bittercress	<i>Solanum americanum</i> <i>Cardamine</i> sp.	Evening primrose, common Evening primrose, cutleaf Flaree, redstem	<i>Oenothera biennis</i> <i>Oenothera laciniata</i> <i>Erodium cicutarium</i>	Plantain, Paleseed Prostrate knotweed Prostrate spurge	<i>Plantago virginica</i> <i>Polygonum aviculare</i> <i>Euphorbia maculata</i>
California burclover Canada thistle, Common (seedlings)	<i>Medicago polymorpha</i> <i>Cirsium arvense</i>	Flabane, blackleaved Florida pusley Gromwell, Yellow	<i>Conyza bonariensis</i> <i>Richardia scabra</i> <i>Amsinckia calycina</i>	Purslane, common Ragweed, common Red tasselflower	<i>Portulaca oleracea</i> <i>Ambrosia artemisiifolia</i> <i>Emilia sonchifolia</i>
Chickweed, Common Chickweed, Mouse-ear Clover, White Corn speedwell	<i>Stellaria media</i> <i>Cerastium vulgatum</i> <i>Trifolium repens</i> <i>Veronica arvensis</i>	Groundsel, common Hairy flabane Hairy nightshade	<i>Senecio vulgaris</i> <i>Erigeron bonariensis</i> <i>Solanum sarcocoides</i>	Redroot pigweed Redstem flabane/Storksbill Russian Thistle	<i>Calandrinia caulescens</i> <i>Amaranthus retroflexus</i> <i>Erodium cicutarium</i>
Cudweed, Linear-leaf/purple Curly dock (seedlings) Cutleaf evening primrose Dandelion, cat's ear Dandelion, common (seedlings)	<i>Gnaphalium purpureum</i> <i>Rumex crispus</i> <i>Oenothera laciniata</i> <i>Hypochaeris radicata</i> <i>Taraxacum officinale</i>	Horseweed/ Marestalk Kochia Lambsquarters, common Lawn burweed	<i>Erigeron canadensis</i> <i>Kochia scoparia</i> <i>Chenopodium album</i> <i>Soliva sessilis</i>	Shepherd's-purse Sowthistle, Annual Spotted catsear Swincress	<i>Capsella bursa-pastoris</i> <i>Sonchus oleraceus</i> <i>Hypochaeris radica</i> <i>Coronopus didymus</i>
Doveweed Eclipta	<i>Murdannia nudiflora</i> <i>Eclipta alba</i>	Long-stalk phyllanthus Panicle willowweed Plantain, Buckhorn	<i>Malva parviflora</i> <i>Phyllanthus tenellus</i> <i>Epilobium paniculatum</i> <i>Plantago lanceolata</i>	Tropic ageratum Velvetleaf Wild buckweed (seedlings) Wild mustard	<i>Ageratum conyzoides</i> <i>Abutilon theophrasti</i> <i>Polygonum convolvulus</i> <i>Sinapis arvensis</i>

Grasses and Sedges Controlled

Annual bluegrass Barnyardgrass, common Cheatgrass Crabgrass Crabgrass, Henry Crabgrass, Large/Hairy Crabgrass, Smooth	<i>Poa annua</i> <i>Echinochloa crus-galli</i> <i>Bromus secalinus</i> <i>Digitaria</i> sp <i>Digitaria ciliaris</i> <i>Digitaria sanguinalis</i> <i>Digitaria ischaemum</i>	Foxtail bromo Foxtail, Giant Foxtail, Green Foxtail, Yellow Goosegrass Guineagrass Mouse barley	<i>Bromus rubens</i> <i>Setaria faberi</i> <i>Setaria viridis</i> <i>Pennisetum glaucum</i> <i>Eleusine indica</i> <i>Panicum maximum</i> <i>Hordeum murinum</i>	Rice flatsedge Ryegrass, Italian Ryegrass, Perennial Sedge, Annual Sprangletop Tufted lovegrass	<i>Cyperus iria</i> <i>Lolium multiflorum</i> <i>Lolium perenne</i> <i>Cyperus compressus</i> <i>Leptochloa</i> spp. <i>Eragrostis pectinacea</i>
---	--	---	--	--	--

(continued)

Weeds Controlled or Suppressed by Esplanade 200SC (continued)					
Weeds Suppressed					
Black medic	<i>Medicago lupulina</i>	Sesbania, hemp	<i>Sesbania exaltata</i>	Wild carrot	<i>Daucus carota</i>
Black mustard	<i>Brassica nigra</i>	Sida, prickly/teaweed	<i>Sida spinosa</i>	Woodsorrel, yellow	<i>Oxalis stricta</i>
False chamomile	<i>Matricaria maritima</i>	Southern brassbuttons	<i>Cotula australis</i>	Woodsorrel/Oxalis	<i>Oxalis species</i>
London rocket	<i>Sisymbrium irio</i>	Sunflower, common	<i>Helianthus sp.</i>		
Prickly lettuce	<i>Lactuca serriola</i>	Vetch, purple	<i>Vicia benghalensis</i>		

Use Sites for Esplanade 200 SC	Rate Range (fl oz/A)	Maximum Single Use Rate (fl oz/A)	Maximum Total Yearly Rate (fl oz/A)
Rail and Rail Yards	5-7	7	10
Managed Roadsides	5-7	7	10
Warm Season Turf Release	3.5-5	5	10
All other use sites	5-7	7	10

Bareground Applications for Non-Residential Non-Crop Sites

Bareground is desired at many non-crop sites for reducing fire hazards, maintaining appropriate lines-of-site, and aesthetic considerations. Examples of sites include but are not limited to guardrails and some median strips near highways, hardscapes, parks, airports, utilities, government and military installations, around farm buildings, plant sites, educational facilities, parking lots, and managed areas. Esplanade 200 SC may be used alone for residual weed control or in tank mixture. Tank mixtures with post emergent herbicides help to control existing weeds. Observe use restrictions for all herbicides if a tank mixture is applied. Use-rates for bareground applications depend on the duration of weed control desired and the weed species listed on this label. Apply Esplanade 200 SC at 5-7 fl oz per acre. A repeat application can be made but not to exceed a total amount of 10 fl oz per acre per year.

Restriction: Applications to hardscapes (e.g. patios, paved parking lots and walkways) may be made by spot application only.

Railroads and Rail Yards

Esplanade 200 SC may be used for preemergent residual control of certain weeds near railroad tracks, ballasts, and rail yards. Follow application instructions under **Bareground Applications** where bareground is the desired result. In situations where warm season turfgrass coverage is desired, such as at railroad crossings, follow use directions under the **Warm Season Turf Release** section of this label. Apply Esplanade 200 SC at 5-7 fl oz per acre. A repeat application can be made but not to exceed a total amount of 10 fl oz per acre per year.

Warm Season Turf Release

Esplanade 200 SC may be used to promote the growth of warm season grasses in areas where low maintenance vegetation or erosion control is desired. Established bermudagrass (*Cynodon dactylon*), centipedegrass (*Eremochloa ophiuroides*), bahiagrass (*Paspalum notatum*) and buffalograss (*Buchloe dactyloides*) are tolerant to Esplanade 200 SC at rates up to 5 fl oz per acre. Application of Esplanade 200 SC in the spring or fall to these grasses will control labeled weeds and allow low maintenance turf to develop. A repeat application can be made but not to exceed a total amount of 10 fl oz per acre per year. Cool season grasses such as Kentucky bluegrass (*Poa pratensis*), perennial ryegrass (*Lolium perenne*), and fescues (*Festuca* sp.) are not tolerant to Esplanade 200 SC. Use Esplanade 200 SC on these grasses only when removal of these grasses is desired.

Esplanade 200 SC can inhibit the emergence of seed and damage newly emerged seedlings. Seeding into turf treated with Esplanade 200 SC should be delayed until at least 8 months after application. Applications to newly seeded turf made sooner than 8 months after emergence may significantly reduce stand establishment and turf vigor.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

Bayer (reg'd), the Bayer Cross (reg'd), Finale®, Esplanade™ and Backed by Bayer™ are trademarks of Bayer.