

**Herbicide** 

New York State Department of Environmental Conservation Division of Materials Management Pesticide Product Registration

**ACCEPTED** 

FOR REGISTRATION

May 16, 2012

Doc id: 531469

**Net Weight:** 

EPA Reg. No. 228-654 EPA Est. No. 39578-TX-001

Manufactured for Nufarm Americas Inc. 150 Harvester Drive Burr Ridge, IL 60527

Nufarm

A broad-spectrum dispersible granule herbicide for use on specified non-crop sites and under paved areas.

62.22%

30.00%

.100.00%

# TOTAL: KEEP OUT OF REACH OF CHILDREN CAUTION/ PRECAUCION

Imazapyr(2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid).

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION/PRECAUCIÓN

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are: barrier laminate, butyl rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, or viton  $\geq 14$ mils. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart. All pilots must wear:

- All pliots must wear:

   Long sleeved shirt and long pants, and

   Shoes, plus socks.

  All mixers, loaders, other applicators and handlers must wear:

   Long sleeved shirt and long pants,

   Shoes plus socks,

   Chospiel createst elegen.

Diuron (3-[3,4-dichloropheny]-1,1-dimethylurea)...

- Chemical-resistant gloves
- with approval number prefix TC-21C,

   Chemical resistant apron when mixing, loading, or cleaning equipment or spills.

See Engineering Controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them.

NIOSH approved dust/mist filtering respirator equipped with any N2 or R, P or HE filter media or an NIOSH-approved dust/mist filtering respirator

## ENGINEERING CONTROLS

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

Flaggers supporting aerial applications must use an enclosed cab that meets the definition in the Worker Protection Standard for Agricultural Pesticides [40 CFR 170.240(d)(5)] for dermal protection. In addition, flaggers must wear long-sleeved shirt, long pants, shoes, and socks.

## USER SAFETY RECOMMENDATIONS

- was should.

  Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

  Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

  Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and cleaned the control of the control

	FIRST AID					
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.     Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.     Call a poison control center or doctor for treatment advice.					
IF ON SKIN OR CLOTHING	Take off contaminated clothing.     Rinse skin immediately with plenty of water for 15 to 20 minutes.     Call a poison control center or doctor for treatment advice.					

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

# ENVIRONMENTAL HAZARDS

ENVIRONMENTAL HAZARDS

This product is toxic to plants. Drift and run-off may be hazardous to plants in water adjacent to treated areas. For terrestrial uses. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS
Spray solutions of this product should be mixed, stored and applied only in stainless steel, fiberglass, plastic, and plastic-lined steel containers.
DO NOT mix, store or apply this product or spray solutions of this product in unlined steel (except stainless steel) containers or spray tanks.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

USE RESTRICTIONS

DO NOT enter or allow others to enter treated areas until sprays have dried.

- Aerial application is prohibited except for application to rights-of-way. . DO NOT USE IN CALIFORNIA.
- DO NOT apply more than a total of 12 lbs. a.i. of diuron (19 pounds per acre of this product) or more than two applications of diuron in a 12-
- monin period.

  Do NoT apply more than 12 lbs. ai/A of diuron per application in areas of high rainfall or dense vegetation. Do not apply more than 8 lbs. ai/A of diuron per application in all other areas.

  Do NoT reapply his product or any other product containing diuron within 90 days of treatment with any product containing diuron.

  Do NoT mix, store or apply this product or spray solutions of this product in unlined steel (except stainless steel) containers or spray tanks.

  DO NOT use on food or feed crops.

  DO NOT treating trigation diffuse or waste used for coop irrigation or for demostic numbers.
- **DO NOT** treat irrigation ditches, or water used for crop irrigation or for domestic purposes
- DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.
- DO NOT contaminate water when disposing of equipment washwaters.
   DO NOT drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where
- the chemical may be washed or moved into contact with their roots.

- the chemical may be washed or moved into contact with their roots.

   DO NOT use on turfgrass at residential sites (including homes, apartment complexes, condominium grounds, daycare facilities, schools, playgrounds, parks, recreational areas, and sports fields).

   DO NOT side trim desirable vegetation with this product.

   DO NOT allow sprays to drift on to desirable plants.

   DO NOT allow sprays to drift on to desirable plants.

   Be sure to clean application equipment after using this product by thoroughly flushing with water.

   DO NOT apply this product with a spoon, a pump-feed backpack spreader or a gravity feed backpack spreader.

   Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for Injunctive relief in Washington Toxics Coalition, et.al. v. EP. C01-0132C, Wb.D. WA). For further information, please refer to <a href="http://www.epa.gov/espp/wtc/">http://www.epa.gov/espp/wtc/</a>.

This product is a dispersible granule that is mixed with water and a spray adjuvant and applied as a spray solution to the following noncropland areas where bare ground is desired: industrial non-crop areas including utility plant sites, petroleum tank farms, pumping installations, storage areas, railroads, utility, and pipeline rights-of-way; highway rights-of-way; non-irrigation ditchbanks; fence rows; farmyards; and non-crop areas around farm buildings. This product may also be used to control weeds under paved surfaces.

This product controls most annual and perennial grasses and broadleaf weeds in addition to many brush and vine species. This product also provides residual control of weeds that germinate in treated areas.

For annual weed control, either preemergence or postemergence applications may be used; however, a late preemergence to early postemergence application provides the best results in most situations.

For perennial weed control, this product is only effective when applied postemergence and will not control perennial weeds that have not emerged at the time of application. For best results, applications should be made when the weeds are growing vigorously and the spray solution should include a spray adjuvant. For specific instructions, see the "Adjuvants" section of this label.

The duration of residual weed control depends upon the types of weeds present, the application rate, and weather conditions. Longer residual control

occurs in areas with sensitive weed species, higher product use rates, lower precipitation and cooler soil temperatures. Higher than average rainfall or warmer than normal temperatures can significantly affect the residual control this product provides and shorten the overall length of control.

Precautions for Avoiding Injury to Non-Target Plants
Untreated trees may be affected by root uptake of this product through movement into the topsoil and injury or loss of desirable trees or other plants may result if this product is applied on or near desirable trees or other plants, on areas where their roots extend, or in locations where the roots extend, or in locations where their roots extend to it may be unabled or may be unabled or the root set in the root set in the root set when the root is all when the root level. We have the root set in the root set when the root is all when the root is all when the root is all the roots and the root set when the root is all the roots and the root set when the root is all the roots and the root set when the root is all the roots and the root set when the root set wh treated soil may be washed or moved into contact with their roots. Treatment of powdery dry soil or light sandy soil when there is little likelihood or rainfall soon after treatment may result in off target movement and possible damage to desirable plants when soil particles are moved by water and/or wind. Exposure to this product may injure or kill most crops and rijury to crops may result if treated soil washed, blown or moved onto land used to produce crops.

## SPRAY DRIFT

Use best practices to avoid drift to all other crops and non-target areas. **DO NOT** apply when conditions favor drift from target areas. The interaction of many equipment- and weather-related factors determine the potential for spray drift. Avoiding spray drift at the application site is the responsibility of the applicator. The applicator must follow the most restrictive precautions to avoid drift, including those found in the ableting as well as applicable state and local regulations and ordinances. A drift control agent may reduce drift, however, it may also decrease weed control. Aerial Applications:

- Action Applications are required to use a Coarse or coarser droplet size (ASABE S572) or, if specifically using a spinning atomizer pozzle, applicators are required. A polication and unique to the dead of the effects of nozel or including an engine of the effects of nozel or interest. Applicators are required to use a Vent of the effects of nozel or interest or interest of the effects of nozel or interest or interest of the effects of nozel or interest or
- neglist above to deet, Application state consider the friends of moze of minimum and might speed when deeminiming displacement.

  (3) The spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The boom length must not exceed 60% of the wingspan or 90% of the rotro blade dismeter to reduce spray drift.

  (4) Applications with wind speeds less than 3 mph and with wind speeds greater than 10 mph are prohibited.
- (5) Applications into temperature inversions are prohibited.(6) **DO NOT** apply by air if sensitive non-target crops are within 100 feet of the application site.

Ground Boom Applications:

Apply with nozzle height no more than 4 feet above the ground or plant canopy and Coarse or coarser droplet size (ASABE S572) or, if specifically using a spinning atomizer nozzle, applicators are required to use a volume mean diameter (VMD) of 385 microns or greater. Use the lowest nozzle height consistent with safety and efficacy.

Apply only when wind speed is less than or equal to 10 miles per hour

## DO NOT apply into temperature inversions. APPLICATION INSTRUCTIONS

For rights-of-way and non-crop areas: The maximum rate per application is 19 lbs./acre of this product (equivalent to 12 lbs. diuron active ingredient per acre) in areas of high rainfall or dense vegetation.

- For all other areas, the maximum rate per application is 13 lbs./acre of this product (equivalent to 8 lbs. diuron active ingredient per acre)
- Make a maximum of two applications per year

The minimum retreatment interval is 90 days

Mix this product in water and apply the specified gallons per acre of spray volume using properly calibrated equipment to deliver a uniformly distributed spray pattern. Apply this product at 7-19 pounds of product per acre, although rates as low as 5 pounds per acre may be used only if tank mixed with another herbicide (see the TANK MIXES section below). For retreatment purposes within the same growing season, apply this product at a rate of less than 7 pounds per acre. Use sufficient volume to insure thorough coverage.

Rainfall may significantly affect length of residual weed control achieved with this product and in cases of increasing rainfall amounts, higher rates may need to be applied to achieve the desired residual control. Refer to the following table for product rates for different amounts rainfall amounts. Actual use rates will depend upon the length of residual control desired as well as weed pressure and environmental conditions.

Average Annual Rainfall	Product Rate	
Less than 15 inches	7-10 pounds†	
Between 15 and 35 inches	8-13 pounds	
Greater than 35 inches	13-19 pounds	

† Initial applications of this product may be made at rates as low as 5-6 pounds per acre, but must be tank mixed with another herbicide (see the

When both mixing and spraying, be sure to maintain sufficient agitation to keep product suspended in spray mixture. Postemergence Applications. When making postemergence applications, always use a spray adjuvant (see "Adjuvants" section of this label) For best results on tough to control perennial weeds, applications should be made in combination with one quart per acre of methylated seed oil Use sufficient volume to insure thorough coverage. For faster burndown or brown-out of target weeds, tank mix this product with products such as Razor®, Razor Pro®, Credit®, or Finale® (refer to the TANK MIXES section for specific instructions).

Tank Mixes

Tank M

NOTE: Tank-mixes with 2,4-D or products that contain 2,4-D have resulted in reduced perennial weed control performance

For Control of Undesirable Weeds under Paved Surfaces

Apply this product only to areas that have been prepared according to good construction practices. All rhizomes, stolons, tubers, or other vegetative plant parts present in the site should be removed by scalping with a grader blade to a depth sufficient to ensure their complete removal. Apply this product under asphalt, pond liners and other paved areas only in industrial sites or where the pavement has a barrier along the perimeter that will prevent encroachment of roots of desirable plants. Sites should be paved as soon as possible after application.

NOTE: DO NOT use this product where landscape plantings could be anticipated, or under paved areas such as driveways or parking lots on residential properties. DO NOT use in recreational areas such as under bike or jogging paths, golf cart paths, or tennis courts. Injury or death of desirable plants may result if this product is applied where roots are present or where they may extend into the treated area.

Application Directions for Paved Surfaces:
When final grade is established, apply this product in sufficient water to ensure thorough and uniform wetting of the soil surface, including the shoulder areas. **DO NOT** move soil following application. In the spray tank, mix clean water with 19 pounds of product per acre during the filling

should areas. Do not inversion forming application, in the spay alian, this dear water with 15 pounds of product per acte utiling the initial population and be sure to agitate before spraying. Incorporation of this product is required to activate the herbicide if the soil is not moist prior to treatment. Use a rototiller or disc to incorporate product into the soil to a depth of 4 to 6 inches. One inch of rainfall or irrigation will also provide uniform incorporation. If using water to incorporate, do not allow treated soil to wash or move into untreated areas.

Adjuvants

Adjuvants glycol should not be considered as surfactants to meet the above requirements).

glycol should not be considered as surfactants to meet the above requirements).

Methylated Seed Oils or Vegetable Oil Concentrates: Methylated seed oils are the adjuvant of choice and research indicates that these oils may aid in the deposition and uptake of this product by plants under moisture or temperature stress and will increase control of perennial weeds. Use a methylated seed oil or vegetable-based seed oil concentrate at the rate of 1.5 - 2 pints per acre in place of a surfactant. When using spray volumes greater than 30 gallons per acre, methylated seed oil or vegetable based seed oil concentrates should be mixed at a rate of 1% of the total spray volume, or a nonionic surfactant as described above may be used instead.

Silicone-Based Surfactants: Silicone-based surfactants may allow greater spreading on the leaf surface as compared to conventional nonionic surfactants by reducing the surface tension of the spray droplets. However, some silicone-based surfactants may limit herbicide uptake by drying too rapidly. Refer to the manufacturer's label for specific rate instructions.

Fertilizer/Surfactant Blends: Use 2 - 3 pints of nitrogen-based liquid fertilizers (such as 28%N, 32%N, 10-34-0, or ammonium sulfate) per acre in combination with the specified rate of nonionic surfactant, methylated seed oil or vegetable/seed oil concentrate. Do NOT use fertilizers in a tank mix without a nonionic surfactant, methylated seed oil or vegetable/seed oil concentrate.

Weeds Controlled

Weeds Controlled

Weeds Controlled

When used at the rates listed in the APPLICATION INSTRUCTIONS section, this product provides preemergence or postemergence control with residual control (control of newly germinating seedlings) of the following target vegetation species. In general, preemergence and postemergence applications of this product control annual weeds while postemergence applications control established biennials and perennials. Use this product control menual product control annual weeds while postemergence applications control established biennials and perennials. Use this product

applications of this product control annual weeds while postemergence applications control established blennials and perennials. Use this product only in accordance with the instructions on this label.

Note Regarding Resistant Biotypes: Naturally occurring biotypes of some of the weeds listed on this label (pigweed, kochia and Russian thistle for example) may not be effectively controlled by this and/or other herbicides with the ALS/AHAS enzyme inhibiting mode of action (such as OUST). To ensure control if naturally occurring ALS/AHAS resistant biotypes are present in an area, tank mix or apply this product sequentially with an appropriate registered herbicide having a different mode of action.

## WEEDS CONTROLLED<sup>1</sup> Growth

Habit <sup>2</sup>	Name (Species)	Habit <sup>2</sup>
A	Kyllinga (Cyperus brevifolius)	A
A	Lovegrass (Eragrostis spp.)	A/P
A	Maidencane (Arundinaria amabilis)	P
P	Orchardgrass (Dactylis glomerata)	P
A	Paragrass (Brachiaria mutica)	P
P	Peppergrass (Lepidium virginicum)	A
P	Phragmites (Phragmites australis)	P
P	Prairie cordgrass (Spartina pectinata)	P
A	Prairie threeawn (Aristida oligantha)	P
P	Quackgrass (Agropyron repens)	P
P	Rattail fescue (Vulpia myuros)	A
A	Reed canarygrass (Phalaris arundinacea)	P
P	Ricegrass (Oryzopsis hymenoides)	A
A	Saltgrass 7.8.9 (Distichlis stricta)	P
P	Sand dropseed 7 (Sporobolus cryptandrus)	P
A	Sandbur (Cenchrus spp.)	A
A	Smooth brome (Bromus inermis)	P
P	Sprangletop 6,7 (Leptochloa spp.)	A
A/P	Timothy (Phleum pretense)	P
A	Torpedograss (Panicum repens)	P
A	Vaseygrass (Paspalum urvillei)	P
P	Velvetgrass (Holcus lanatus)	A
A	Wild barley (Hordeum spp.)	A
P	Wild oats (Avena fatua)	A
P	Wirestem muhly (Muhlenbergia frondosa)	P
	Witchgrass (Panicum capillare)	A
	A A A P P A A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A A P P A A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A A P P A A A A A P P A A A A A P P A A A A A P P A A A A P P A A A A P P A A A A P P A A A A P P A A A A A P P A A A A A P P A A A A A P P A A A A A P P A A A A A P P A A A A A P P A A A A A P P A A A A A P P A A A A A P P A A A A A P P A A A A A P P A A A A A P P A A A A A P P A A A A A A P P A A A A A P P A A A A A A P P A A A A A P P A A A A A P P A A A A A A P P A A A A A A P P A A A A A A P P A A A A A A A A A A A A A A A A A A A A	A Kyllinga (Cyperus brevifolius) A Lovegrass (Fragrostis spp.) A Maidencane (Arundinaria amabilis) P Orchardgrass (Dactylis glomerata) A Paragrass (Dactylis glomerata) A Paragrass (Brachiaria mutica) P Peppergrass (Lepidium virginicum) P Phragmites (Phragmites australis) P Prairie cordgrass (Spartina pectinata) A Prairie threeawn (Aristida oligantha) P Quackgrass (Agropyron repens) P Rattali fescue (Vulpia myuros) P Rattali fescue (Vulpia myuros) A Reed canarygrass (Phalaris arundinacea) P Ricegrass (Oryzopsis hymenoides) A Saltgrass <sup>7,80</sup> (Distichiis stricta) P Sand dropseed <sup>71</sup> (Sporobolus cryptandrus) A Sanodhur (Cenchrus spp.) A Sanothor (Cenchrus spp.) A Smooth brome (Bromus inermis) P Sprangletop <sup>52</sup> (Leptochloa spp.) A/P Timothy (Phieum pretense) A Torpedograss (Paspalum urville) P Velvetgrass (Holcus lanatus) A Wild barley (Hordeum spp.) P Wilf oats (Avena fatua) P Wilf estem mulhy (Muhlenbergia frondosa)

		Witchgrass (Panicum capillare)	A
BROADLEAF WEEDS Name (Species)	Growth Habit <sup>2</sup>	Name (Species)	Growth Habit <sup>2</sup>
Arrowwood (Pluchea sericea)	A	Nettleleaf goosefoot (Chenopodium murale)	A
Ageratum (Asteraceae houstonianum)	P	Oxeye daisy (Chrysanthemum leucanthemum)	P
Broom snakeweed 3 (Gutierrezia sarothrae)	P	Pennycress (Thlaspi spp.)	A
Bull thistle (Cirsium vulgare)	В	Pepperweed (Lepidium spp.)	A
Burdock (Arctium spp.)	В	Pigweed <sup>6</sup> (Amaranthus spp.)	A
Canada thistle7 (Cirsium arvense)	P	Pineapple weed (Matricaria matricarioides)	P
Carolina geranium (Geranium carolinianum)	A	Plantain (Plantago spp.)	P
Carpetweed (Mollugo verticillata)	A	Pokeweed (Phytolacca Americana)	P
Clover (Trifolium spp.)	A/P	Prickly sida (Sida spinosa)	A
Cocklebur (Xanthium strumarium)	A	Primrose (Oenothera kunthiana)	P
Common chickweed (Stellaria media)	A	Puncturevine (Tribulus terrestris)	A
Common ragweed (Ambrosia artemisiifolia)	A	Purple loosestrife 3 (Lythrum salicaria)	P
Corn spurry (Spergula arvensis)	P	Purslane (Portulaca spp.)	A
Dandelion (Taraxacum officinale)	P	Ragweed (Ambrosia spp.)	A
			continue

continued **BROADLEAF WEEDS** Name (Species) Name (Species) Δ/P Rush skeletonweed 3 (Chondrilla juncea Desert Camelthorn (Alhagi pse Russian knapweed (Centaurea repens Dock (Rumex spp.) Saltbush (Atriplex spp.) Dogfennel (Eupatorium capillifolium Sesbania (Sesbania spp.) ilaree (*Erodium* spp.) Fleahane (Frigeron snn.) Silverleaf nightshade (Solanum elaeagnifolium) Giant ragweed 7 (Ambrosia trifida) Sheperd's-purse (Capsella bursa-pastoris) Smartweed (Polygonum spp.) A/P Goldenrod (Solidago spp.) Grey rabbitbrush (Chrysothamnus nauseosus) Sorrell (Rumex spp.) A A/P roundcherry (*Physalis* spp.) Sneedwell (Veronica spp.) Α Hawksbeard (Crepis spp.) Stinging nettle <sup>3</sup> (*Urtica dioica*)
Sunflower (*Helianthus* spp.) A/B Horsenettle (Solanum Canadensis Sweet clover (Melilotus spp.) Indian mustard (Brassica juncea) Texas thistle (Cirsium texanum) Α Japanese bamboo (Polygonum cuspisatum) Velvetleaf (Abutilon theophrasti Western ragweed (Ambrosia psilostachya) nawel (Scleranthus annuus Kochia 3 (Kochia scoparia) Wild buckwheat (Polyonum convolvulus) Α Wild carrot (Daucus carota) Lambsquarters (*Chenopodium album*) Wild lettuce (Lactuca spp.) A/B espedeza (*Lespedeza* spp. Little mallow (Malva parviflora Wild parsnip (Pastinaca sativa) Wild radish (Raphanus raphanistr larigold (*Tagetes* spp.) B B Wild turnin (Brassica camnestris) Milkweed (Asclenias spn Miners lettuce (Montia perfoliata spp.) Woolly leaf bursage (Franseria tomentosa) Morningglory (Ipomoea spp.) Yellow starthistle (Centaurea solstitialis) A P Mullein (Verbascum spp.) Yellow woodsorrel (Oxalis stricta) VINES AND BRAMBLES Growth Habit<sup>2</sup> Blackberry 4 (Rubus spp.) Morningglory (Inomoea spp.) A/P Dewberry 4 (Rubus spp.) Poison Ivy (Rhus radicans) Field bindweed (Convolvulus arvensis Redvine (Brunnichia cirrhosa Greenbriar (Smilax spp.) Trumpetcreeper 7 (Campsis radicans Honeysuckle (Lonicera spp.) Wild buckwheat (Polygonum convolvulus) Kudzu 5 (Pueraria lobata) Wild grape (Vitis spp.)

his product controls over 30 species of brush

Name (Species)	Growth Habit <sup>2</sup>	Name (Species)	Growth Habit <sup>2</sup>
Alder (Alnus spp.)	P	Mulberry (Morus spp.)	P
American beech (Fagus grandifolia)	P	Oak (Quercus spp.)	P
Ash (Fraxinus spp.)	P	Permission (Diospyros virginiana)	P
Bald cypress (Taxodium distichum)	P	Pine 10 (Pinus spp.)	P
Bigleaf maple (Acer macrophyllum)	P	Poplar (Populus spp.)	P
Black locust 10 (Robinia pseudoacacia)	P	Privet (Ligustrum vulgare)	P
Black gum (Nyssa sylvatica)	P	Red alder (Alnus rubra)	P
Boxelder (Acer negundo)	P	Red maple (Acer rubrum)	P
Cherry (Prunus spp.)	P	Russian olive (Elaeagnus angustifolia)	P
Chinaberry (Melia azedarach)	P	Sassafras (Sassafras albidum)	P
Dogwood (Cornus spp.)	P	Sourwood (Oxydendrum arboreum)	P
Elm 11 (Ulmus spp.)	P	Sweetgum (Liquidambar styraciflua)	P
Hawthorn (Crataegus spp.)	P	Water Willow (Justicia americana)	P
Hickory (Carya spp.)	P	Willow (Salix spp.)	P
Honeylocust 10 (Gleditsia triacanthos)	P	Yellow poplar (Liriodendron tulipifera)	P
Maple (Acer spp.)	P		P

Where heavy or well-established infestations occur, use the higher specified rates.

whele heavy or well-examinate microsolists count, use me injure specimen raies.

Growth Habit: A-munal, B-- Biennial, P- Perennial results.

Early postemergence applications are required for best results.

The degree of control is species dependent; some *Rubus* species may not be completely controlled.

Use a minimum of 75 GPA; repeat applications may be required to control established stands.

Control is species dependent, for preemergence control a tank-mix with Pendulum herbicide and/or a postemergence application of a labeled herbicide may be required.

Control is species dependent; for preemergence control a tank-nerbicide may be required.
 A minimum of 13 pounds of this product per acre is required.
 Tank-mix with Oust\*, Spyder\* or Spyder\* Extra for best results.
 Tank mix with glyphosate or triclopyr.
 Tank mix with glyphosate.

Growth

# STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry area in original container

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on-site or at an approved waste disposal facility. CONTAINER HANDLING: Nonrefillable container. DO NOT reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. To The EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER MAD (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS ON THE PROBLEM THE PROPERTY OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR TRADE USAGE. UNINTENDED CONSCIOUENCES, INCLUDING BUT NOT LIMITED TO INFERENTIANE OF USER OF SUCH FACTORS AS THE PRESENCE OR ASSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SET OR PROPERTY OF MAINEY AND ASSIMER OF SURVINE OR USER THIS WEITHER CONTROL OF MANUFACTIONS AND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR

CONSCOURTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DOWN DAMAGES OR SUNS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISINGS OUT OF THE MISUSE, OR FALLIBET TO FOLLOW LASEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

Nufarm Imazuron Herbicide is not manufactured or distributed by BASF, seller of Sahara® Credit®, Spyder® Extra, Tahoe®, Relegate®, Diablo®, Polaris® are Registered Trademarks of Nufarm Americas Inc. Arsenal®, Banvel®, Pendulum®, Plateau® and Sahara® are trademarks of BASF Corporatio Arsenare, Barryere, reflouenine, Fraceau and Garard C Finale® is a trademark of Bayer. Garlon® is a trademark of Dow AgroSciences Company.

Karmex® and Oust® are trademarks of E.I Du Pont de Nemours and Company. Roundup® is a trademark of Monsanto Company.

Vanguish® is a trademark of a Syngenta Group Company

13502000



